

Dressed scalar propagator in a non-Abelian background from the worldline formalism

Naser Ahmadinia^{1,2,*} Fiorenzo Bastianelli^{3,4,†} and Olindo Corradini^{1,5,4,‡}

¹*Facultad de Ciencias en Física y Matemáticas, Universidad Autónoma de Chiapas, Ciudad Universitaria, Tuxtla Gutiérrez 29050, México*

²*Center for Relativistic Laser Science, Institute for Basic Science, Gwangju 61005, Korea*

³*Dipartimento di Fisica ed Astronomia, Università di Bologna, Via Irnerio 46, I-40126 Bologna, Italy*

⁴*INFN, Sezione di Bologna, Via Irnerio 46, I-40126 Bologna, Italy*

⁵*Dipartimento di Scienze Fisiche, Informatiche e Matematiche,*

Università degli Studi di Modena e Reggio Emilia, Via Campi 213/A, I-41125 Modena, Italy

(Received 7 September 2015; published 28 January 2016; corrected 1 February 2016)

We study the propagator of a colored scalar particle in the background of a non-Abelian gauge field using the worldline formalism. It is obtained by considering the open worldline of a scalar particle with extra degrees of freedom needed to take into account the color charge of the particle, which we choose to be in the fundamental representation of the gauge group. Specializing the external gauge field to be given by a sum of plane waves, i.e. a sum of external gluons, we produce a master formula for the scalar propagator with an arbitrary number of gluons directly attached to the scalar line, akin to similar formulas derived in the literature for the case of the scalar particle performing a loop. Our worldline description produces at the same time the situation in which the particle has a color charge given by an arbitrarily chosen symmetric or antisymmetric tensor product of the fundamental.

DOI: [10.1103/PhysRevD.93.025035](https://doi.org/10.1103/PhysRevD.93.025035)

I. INTRODUCTION

The worldline representation of effective actions has seen a great deal of activity in the past twenty years, starting with the work of Strassler [1], who rederived the Bern-Kosower master formulas [2,3] directly from point particle path integrals; see Ref. [4] for a review. Since then, many extensions and applications of the worldline formalism have been considered: multiloop computations [5], non-perturbative worldline methods [6–8], the numerical worldline approach to the Casimir effect [9], the worldline formalism in curved spacetime [10–12], photon-graviton mixing at one loop [13], higher-spin field theory [14–16], the world-graph approach to QCD [17], as well as the heat kernel expansion [18–20] and applications to noncommutative QFT [21,22], Standard Model physics [23] and its grand-unified extensions [24], just to name a few.

Unlike effective actions, the worldline representation of dressed propagators is still a relatively unexplored land, though a worldline representation for the dressed propagator of a scalar field coupled to electromagnetism (scalar QED) was proposed long ago by Feynman [25]. It consists of a worldline path integral where the coordinate paths have the topology of a line, and come with Dirichlet boundary conditions. The problem has then been reconsidered in Ref. [26] along the lines of [1–3]. More recently, the dressed

propagator in a scalar field theory has been studied with worldline methods to address the summation of ladder and crossed-ladder diagrams and analyze the emergence of bound states [27]. A full worldline description of dressed QFT propagators would be quite welcome in gauge theories as well. It may allow one to address several different issues, providing a systematic way of computing scattering amplitudes that could be beneficial both at the perturbative and nonperturbative levels. At the perturbative level it may improve on the efficiency of perturbative calculations and give perhaps a better understanding of color/kinematics dualities [28]. At the nonperturbative level it might be useful to address the emergence of bound states, or to study, for instance, the covariance of the Green's functions under a change in the gauge parameter that gives rise to the so-called Landau-Khalatnikov-Fradkin transformations [29,30]. Indeed the worldline formalism has been used recently to extend these transformations, originally derived for the scalar propagator, to an arbitrary n -point function in scalar QED [31]. The search of computational methods for tree-level amplitudes in gauge theories has been quite an active subject in the past decade. In particular, in scalar QCD, techniques based on recursive relations have been successfully found; see e.g. Refs. [32–35].

Here we consider the worldline approach to the propagator for a scalar field coupled to an external non-Abelian gauge field. Similarly to previous worldline treatments of non-Abelian effective actions [36,37], we obtain the path ordering—needed for the gauge covariance of the worldline path integral—through the quantization of suitable

*naser@ifm.umich.mx

†bastianelli@bo.infn.it

‡olindo.corradini@unimore.it

auxiliary variables [38–40]. Their usefulness resides in the fact that they allow us to get rid of the explicit path-ordering prescription. Computationally it is a great advantage, analogous to the replacement of the path-ordered spin factors, present in Feynman’s original proposal for describing a spin-1/2 particle [25], with a Grassmann path integral over fermionic coordinates [41]. However, the auxiliary variables thus introduced must be constrained in order to produce an irreducible representation (irrep) of the gauge group. One possibility is to couple them to a worldline $U(1)$ gauge field, which allows for a Chern-Simons term whose coupling constant is chosen to project onto the desired irrep. This is the method already seen at work in the worldline description of p -forms [12,42], later extended to the treatment of color charges. Furthermore, it allows us to use commuting auxiliary variables which, in absence of the Chern-Simons term, would generate an infinite-dimensional color space.

Below we show how to employ commuting auxiliary variables with a Chern-Simons term to study the propagator of a scalar particle in the background of a non-Abelian gauge field by representing it as a suitable worldline path integral. The non-Abelian charge (i.e. color) of the scalar particle can be arbitrary, though for simplicity we choose it to be in the fundamental representation. The propagator dressed by the external gauge field is (background) gauge covariant. One may then specialize the external gauge field to be given by a sum of plane waves (external gluons) and perform the path integral. This produces a master formula generating the propagator with an arbitrary number of gluons directly attached to it. At fixed number n of gluons, we call the latter “partial n -gluon scalar propagator”: indeed it consists of the scalar propagator with n gluons directly attached to it, while the gluon self-interactions are excluded. As such it is a gauge dependent object. However this dressed propagator is valid off-shell, and it can be used as a building block for higher-loop amplitudes. For example, being valid off-shell, one could use it to construct ladder diagrams with gluon rungs and study the emergence of bound states, as was done in Ref. [27] for the purely scalar case. This would still be a gauge-dependent ladder, but in the bound-state studies maintaining gauge invariance remains a difficult open problem. One may also try to use it on-shell and study tree-level amplitudes, but then the addition of the reducible diagrams with three- and four-gluon vertices will be essential to achieve gauge invariance. We do not wish to discuss this final issue here, though one may hope that a kind of “tree replacement rules”—such as those used in similar one-loop master formulas to generate the missing one-particle-reducible terms (see for example Ref. [4])—might work here as well. Alternatively, one may combine the present treatment of color charges with the world-graph approach of Ref. [17], which is able to generate particle reducible graphs. We leave this to future analysis.

The main point for us in the present paper is to use and exemplify a novel representation of the color degree of freedom. The partial n -gluon scalar propagator is the most natural starting point for testing its usefulness in QCD applications.

A final comment on the Chern-Simons coupling: fine-tuning such coupling allows us to give the scalar particle a color charge corresponding to any arbitrarily chosen symmetric tensor product of the fundamental representation, rather than to the fundamental itself. Replacing the commuting variables with anticommuting ones would also give a similar construction, but with the non-Abelian charge sitting in an arbitrarily chosen antisymmetric tensor product of the fundamental representation, including the fundamental itself. The choice is again made by selecting an appropriate Chern-Simons coupling.

II. TREE-LEVEL AMPLITUDES IN SCALAR QED

As a warm-up exercise, and in order to fix our notation, we review the computation of tree-level amplitudes in scalar QED from a point particle (worldline) path integral on the line, i.e. with Dirichlet boundary conditions.

It is well known, since the seminal work of Feynman [25], that the propagator for a massive charged scalar field coupled to electromagnetism can be obtained from a worldline path integral

$$\langle \phi(x) \bar{\phi}(x') \rangle_A = \int_{x(0)=x'}^{x(1)=x} \frac{DxDe}{\text{Vol Gauge}} e^{-S[x,e;A]}, \quad (2.1)$$

where the particle action in Euclidean signature, is given by (space-time indices are left implicit where not required)

$$S[x, e; A] = \int_0^1 d\tau \left(\frac{1}{2e} \dot{x}^2 + \frac{m^2 e}{2} - iq\dot{x} \cdot A(x) \right), \quad (2.2)$$

with e being the einbein, i.e. the gauge field for the one-dimensional diffeomorphisms. After the gauge fixing $e = 2T$, the path integral over e reduces to a numerical integral over inequivalent constant gauge configurations labeled by the proper time T , i.e.

$$\langle \phi(x) \bar{\phi}(x') \rangle_A = \int_0^\infty dT e^{-Tm^2} \int_{x(0)=x'}^{x(1)=x} Dx e^{-\int_0^1 d\tau \left(\frac{1}{2T} \dot{x}^2 - iq\dot{x} \cdot A(x) \right)}. \quad (2.3)$$

In particular, treating the external electromagnetic potential as a perturbation, the expression (2.3) can be shown to contain the sum of an infinite number of tree-level Feynman diagrams with an incoming scalar particle in x' , an outgoing scalar particle in x , and an arbitrary number of photons. Specifically, in order to extract the amplitude with n photons, one first writes the potential as a sum of the n photons with polarizations ε_i , and momenta k_i ,

$$A_\mu(x(\tau)) = \sum_{l=1}^n \varepsilon_{\mu,l} e^{ik_l \cdot x(\tau)}, \quad (2.4)$$

and expands the exponential involving the potential, then extracts the amplitude as the term in (2.3) that is multilinear in all the different polarizations ε_l 's. The amplitude, thus, reads

$$\begin{aligned} \mathcal{A}(x', x; \varepsilon_1, k_1, \dots, \varepsilon_n, k_n) &= (iq)^n \int_0^\infty dT e^{-Tm^2} \prod_{l=1}^n \int_0^1 d\tau_l \\ &\times \int_{x(0)=x'}^{x(1)=x} Dx e^{-\frac{1}{4T} \int_0^1 d\tau \dot{x}^2} e^{\sum_l (ik_l \cdot x_l + \varepsilon_l \cdot \dot{x}_l)} \Big|_{\text{m.l.}}, \end{aligned} \quad (2.5)$$

where the $\dot{x} \cdot \varepsilon$ terms have been reexponentiated and $x_l := x(\tau_l)$, and where ‘‘m.l.’’ stands for multilinear. We may now perform the path integral by splitting the generic path $x(\tau)$ into the background, $x_{\text{bg}}(\tau) = x' + (x - x')\tau$, satisfying the boundary conditions, and quantum fluctuations $y(\tau)$ with vanishing boundary conditions. We, thus, get

$$\begin{aligned} \mathcal{A}(x', x; \varepsilon_1, k_1, \dots, \varepsilon_n, k_n) &= (iq)^n \int_0^\infty \frac{dT}{(4\pi T)^{\frac{D}{2}}} e^{-Tm^2 - \frac{1}{4T}(x-x')^2} \prod_{l=1}^n \int_0^1 d\tau_l \\ &\times e^{\sum_l [ik_l \cdot (x' + \tau_l(x-x')) + \varepsilon_l \cdot (x-x')]} \langle e^{\sum_l (ik_l \cdot y_l + \varepsilon_l \cdot \dot{y}_l)} \rangle \Big|_{\text{m.l.}}, \end{aligned} \quad (2.6)$$

where the expectation value $\langle \dots \rangle$ is taken with respect to the free Gaussian path integral $\int_{y(0)=0}^{y(1)=0} Dy e^{-\frac{1}{4T} \int_0^1 d\tau \dot{y}^2}$. Expression (2.6) is, thus, the expectation value of the product of photon vertex operators

$$V_A[\varepsilon, k] = e^{ik \cdot x' + \varepsilon \cdot (x-x')} \int_0^1 d\tau e^{[ik \cdot (\tau(x-x') + y) + \varepsilon \cdot \dot{y}]}. \quad (2.7)$$

$$\begin{aligned} \mathcal{A}(p', p; \varepsilon_1, k_1, \dots, \varepsilon_n, k_n) &= \int d^D x \int d^D x' e^{i(p \cdot x + p' \cdot x')} \mathcal{A}(x', x; \varepsilon_1, k_1, \dots, \varepsilon_n, k_n) \\ &= \int d^D x_+ \int d^D x_- e^{i(p+p') \cdot x_+ + \frac{i}{2}(p-p') \cdot x_-} \mathcal{A}\left(x_+ - \frac{x_-}{2}, x_+ + \frac{x_-}{2}; \varepsilon_1, k_1, \dots, \varepsilon_n, k_n\right), \end{aligned} \quad (2.13)$$

with $x_- := x - x'$ and $x_+ := \frac{x+x'}{2}$. The integral over the ‘‘center of mass’’ x_+ yields the energy-momentum conservation delta function $(2\pi)^D \delta^{(D)}(p + p' + \sum k_l)$, whereas the integral over the ‘‘distance’’ x_- is Gaussian. Hence, after some simple manipulations, the amplitude reduces to

$$\begin{aligned} \tilde{\mathcal{A}}(p', p; \varepsilon_1, k_1, \dots, \varepsilon_n, k_n) &= (iq)^n \int_0^\infty dT e^{-T(m^2 + p^2)} \prod_{l=1}^n \int_0^1 d\tau_l \\ &\times \exp\left\{ T(p-p') \cdot \sum_{l=1}^n (-k_l \tau_l + i\varepsilon_l) + T \sum_{l,l'} [k_l \cdot k_{l'} \Delta_{l-l'} - 2i\varepsilon_l \cdot k_{l'} \dot{\Delta}_{l-l'} + \varepsilon_l \cdot \varepsilon_{l'} \ddot{\Delta}_{l-l'}] \right\} \Big|_{\text{m.l.}}, \end{aligned} \quad (2.14)$$

From the free path integral one may obtain the worldline propagator

$$\langle y^\mu(\tau) y^\nu(\tau') \rangle = -2T \delta^{\mu\nu} \Delta(\tau, \tau') \quad (2.8)$$

$$\Delta(\tau, \tau') = \tau\tau' + \frac{1}{2}|\tau - \tau'| - \frac{1}{2}(\tau + \tau'), \quad (2.9)$$

and Eq. (2.6) can, thus, be written as

$$\begin{aligned} \mathcal{A}(x', x; \varepsilon_1, k_1, \dots, \varepsilon_n, k_n) &= (iq)^n \int_0^\infty \frac{dT}{(4\pi T)^{\frac{D}{2}}} e^{-Tm^2 - \frac{1}{4T}(x-x')^2} \prod_{l=1}^n \int_0^1 d\tau_l \\ &\times \exp\left\{ \sum_{l=1}^n [ik_l \cdot (x' + \tau_l(x-x')) + \varepsilon_l \cdot (x-x')] \right. \\ &\left. + T \sum_{l,l'} (k_l \cdot k_{l'} \Delta_{l-l'} - i2\varepsilon_l \cdot k_{l'} \dot{\Delta}_{l-l'} - \varepsilon_l \cdot \varepsilon_{l'} \ddot{\Delta}_{l-l'}) \right\} \Big|_{\text{m.l.}}, \end{aligned} \quad (2.10)$$


where $\Delta_{l-l'} := \Delta(\tau_l, \tau_{l'})$, whereas left and right bullets indicate derivatives with respect to τ_l and $\tau_{l'}$ respectively. Therefore,

$$\bullet \Delta_{l-l'} = \tau_{l'} - \theta(\tau_{l'} - \tau_l) \quad (2.11)$$

$$\bullet \dot{\Delta}_{l-l'} = 1 - \delta(\tau_l - \tau_{l'}), \quad (2.12)$$

where $\theta(x)$ is the Heaviside function. Notice that—although products of delta functions appear in the full expansion of the exponential in the expression (2.10)—the multilinear part of the expansion, that yields the amplitude, does not involve singular, or ill-defined, terms.

In order to get the amplitude fully in momentum space, one may Fourier transform the expression (2.10) and get


 FIG. 1. Diagram structure for the n -photon amplitude.

with $\Delta_{l-l'} := \frac{1}{2}|\tau_l - \tau_{l'}|$, i.e. the translation-invariant linear part of the worldline propagator (2.9), a fact that was already noted in Ref. [26]. For simplicity, and for later convenience, in Eq. (2.14) the overall energy-momentum conservation delta function has been stripped off, and the resulting amplitude has been indicated with $\tilde{\mathcal{A}}$.

Equation (2.14) is the Bern-Kosower-type formula for the tree-level amplitude with n photons and two scalars, originally worked out in Ref. [26] (see also Ref. [31]). Note that this amplitude is amputated on the external photon lines, and may be called the n -photon scalar propagator. It gives the sum of all the QFT Feynman diagrams depicted in Fig. 1, which involve linear vertices and seagull vertices, and where the external photon momenta appear in all possible orderings. This is an advantage of the worldline formalism, that it combines all orderings into a single expression and that, for scalar QFT, generates the seagull vertices within the linear worldline vertex operator. It is easy to check that, for $n = 2$, it reproduces the sum of the three Feynman diagrams responsible for the scalar QED Compton scattering. Notice that the integrand in Eq. (2.14) is made of a worldline translation-invariant double sum, and a simple sum that is not translation-invariant. However it becomes translational invariant on the shell of the scalar lines: indeed, imposing the translation $\delta\tau_l = \lambda$ on such sum, and using momentum conservation, it gives $\lambda(p - p') \cdot \sum_l k_l = \lambda(p'^2 - p^2)$, which vanishes on-shell.

III. DRESSED SCALAR PROPAGATOR IN A NON-ABELIAN BACKGROUND

A straightforward way to generalize the previous results to the case of the coupling of a scalar field to a non-Abelian gauge field $W_\mu(x) = W_\mu^a(x)T^a$, with T^a belonging to the

Lie algebra \mathfrak{g} of a gauge group G , is to use a path-ordering prescription to guarantee the gauge covariance of the path integral. Let us, thus, take a colored scalar field with an index in an irreducible representation of a gauge group, which for definiteness we assume to be $G = SU(N)$. The scalar field propagator in the presence of the non-Abelian gauge field can, thus, be written as

$$\langle \phi_\alpha(x) \bar{\phi}^{\alpha'}(x') \rangle_W = \int_{x(0)=x'}^{x(1)=x} \frac{DxDe}{\text{Vol Gauge}} (\mathcal{P}e^{-S[x,e;W]})_\alpha^{\alpha'}, \quad (3.1)$$

where the particle action in Euclidean signature is given by

$$S[x, e; W] = \int_0^1 d\tau \left(\frac{1}{2e} \dot{x}^2 + \frac{m^2 e}{2} - ig \dot{x} \cdot W(x) \right), \quad (3.2)$$

and with \mathcal{P} denoting the path ordering. Although this expression contains implicitly the multigluon scalar propagator, it may be convenient to investigate an alternative approach that implements the path ordering using some auxiliary variables. For the case of one-loop gluon amplitudes, generated by a scalar field or by a Dirac field, this procedure was proposed in Ref. [40] and more recently reconsidered in Refs. [36,37]. The main advantages of such a procedure are that it is (i) an automatic implementation of path ordering, through the quantization of the auxiliary fields, that allows us to simplify the perturbative calculation and (ii) a simple way to extend our formulas to the case of a scalar field in a generic (anti)symmetric tensor product representation of the gauge group.

For definiteness, let T^a be in the fundamental representation, i.e. $(T^a)_{\alpha}^{\alpha'}$, $\alpha = 1, \dots, N$, then the totally (anti)symmetric tensor $\phi_{\alpha_1 \dots \alpha_r}$ transforms in an irreducible (anti)symmetric product of r fundamental representations. Let us, thus, consider the addition of complex auxiliary variables, c_α and \bar{c}^α , that sit in the fundamental and antifundamental representations of the gauge group, respectively, and that are coupled to the gauge field potential

$$\int_0^1 d\tau [\bar{c}^\alpha \dot{c}_\alpha - ig\dot{x}^\mu W_\mu^a \bar{c}^\alpha (T^a)_\alpha^\beta c_\beta] - \bar{c}^\alpha c_\alpha(1). \quad (3.3)$$

Note the boundary term that allows us to set initial conditions on c and final conditions on \bar{c} . The canonical quantization of such fields gives rise to creation and annihilation operators \hat{c} and \hat{c}^\dagger that generate a suitable Fock space. Using a coherent state basis like the one summarized in Appendix A, it yields a scalar wave function in the coordinate \bar{u} , the left eigenvalue on coherent states of the creation operator c^\dagger , that in turn can be expanded as a \bar{u} -graded sum of fields

$$\phi(x, \bar{u}) = \phi(x) + \bar{u}^\alpha \phi_\alpha(x) + \frac{1}{2!} \bar{u}^{\alpha_1} \bar{u}^{\alpha_2} \phi_{\alpha_1 \alpha_2}(x) + \dots, \quad (3.4)$$

so that if the auxiliary variables are (anti)commuting, the tensors $\phi_{\alpha_1 \dots \alpha_r}$ sit in an (anti)-symmetrized tensor product of r fundamental representations. In the following we will concentrate on the case of commuting auxiliary variables, as the anticommuting counterpart is just a straightforward generalization. Another necessary ingredient is, thus, a projector that allows us to single out from Eq. (3.4) only an irreducible tensor with r indices. This may be achieved by adding a $U(1)$ worldline gauge field to the action, along with a Chern-Simons term. The full locally symmetric action reads

$$S_I[x, c, \bar{c}; e, a, W] = \int_0^1 d\tau \left(\frac{1}{2e} \dot{x}^2 + \frac{m^2 e}{2} - ig\dot{x}^\mu W_\mu^a \bar{c}^\alpha (T^a)_\alpha^\beta c_\beta + \bar{c}^\alpha (\partial_\tau + ia) c_\alpha - isa \right) - \bar{c}^\alpha c_\alpha(1), \quad (3.5)$$

where $a(\tau)$ is the $U(1)$ gauge field and s is the Chern-Simons coupling, to be fixed shortly. Due to the presence of the auxiliary fields, the potential is no longer matrix-valued and no explicit path-ordering is thus needed. The propagator reads

$$\langle \phi(x, \bar{u}) \bar{\phi}(x', u') \rangle_W = \int_{x(0)=x', c(0)=u'}^{x(1)=x, \bar{c}(1)=\bar{u}} \frac{Dx D\bar{c} Dc De Da}{\text{Vol Gauge}} \times e^{-S_I[x, c, \bar{c}, e, a; W]}, \quad (3.6)$$

where u'_α and u_α represent the initial and final color states of the scalar field. Thus, if the scalar field sits in the rank- r totally symmetric representation, the above field operators are (apart from numerical factors) given by

$$\begin{aligned} \phi(x, \bar{u}) &\sim \phi_{\alpha_1 \dots \alpha_r}(x) \bar{u}^{\alpha_1} \dots \bar{u}^{\alpha_r}, \\ \bar{\phi}(x', u') &\sim \bar{\phi}^{\alpha_1 \dots \alpha_r}(x') u'_{\alpha_1} \dots u'_{\alpha_r}. \end{aligned} \quad (3.7)$$

Upon gauge-fixing, the einbein e can be set to a modulus $2T$ as above, whereas the Abelian gauge field can be set to a constant angle modulus $\theta \in [0, 2\pi)$. The gauge-fixed path integral thus reads

$$\begin{aligned} \langle \phi(x, \bar{u}) \bar{\phi}(x', u') \rangle_W &= \int_0^\infty dT e^{-Tm^2} \int_0^{2\pi} \frac{d\theta}{2\pi} e^{is\theta} \int_{x(0)=x'}^{x(1)=x} Dx \\ &\times \int_{c(0)=u'}^{\bar{c}(1)=\bar{u}} D\bar{c} Dc e^{-S[x, c, \bar{c}, 2T, \theta; W]}, \end{aligned} \quad (3.8)$$

with

$$\begin{aligned} S[x, c, \bar{c}; 2T, \theta, W] &= \int_0^1 d\tau \left(\frac{1}{4T} \dot{x}^2 - ig\dot{x}^\mu W_\mu^a \bar{c}^\alpha (T^a)_\alpha^\beta c_\beta \right. \\ &\left. + \bar{c}^\alpha (\partial_\tau + i\theta) c_\alpha \right) - \bar{c}^\alpha c_\alpha(1), \end{aligned} \quad (3.9)$$

where the constant part $-Tm^2 + is\theta$ was stripped off. Classically the equation of motion for a corresponds to the constraint $\bar{c}^\alpha c_\alpha - s = 0$. In the canonical quantization approach, this turns into an operatorial constraint to be imposed *à la* Dirac-Gupta-Bleuler on the wave function. As shown in Appendix A, we choose to identify the path integral expression $c^\dagger c_\alpha$ as the one representing the normal-ordered Hamiltonian $\hat{c}^\dagger \hat{c}_\alpha$. Therefore, the quantum constraint reads

$$\left(\bar{c}^\alpha \frac{\partial}{\partial \bar{c}^\alpha} - s \right) \phi(x, \bar{c}) = 0, \quad (3.10)$$

and thus, in order to single out the tensor with r indices, we identify $s = r$, and set the path integral normalization to [see Eq. (A24)]

$$\int_{c(0)=u'}^{\bar{c}(1)=\bar{u}} D\bar{c} Dc e^{-\int_0^1 d\tau \bar{c}^\alpha (\partial_\tau + i\theta) c_\alpha + \bar{c}^\alpha c_\alpha(1)} = e^{e^{-i\theta} \bar{u} \cdot u'}, \quad (3.11)$$

with $\bar{u} \cdot u' = \bar{u}^\alpha u'_\alpha$. In the following, we find it convenient to twist the auxiliary fields in order to absorb the θ term, i.e. $c(\tau) \rightarrow c(\tau) e^{-i\theta\tau}$ and $\bar{c}(\tau) \rightarrow \bar{c}(\tau) e^{i\theta\tau}$, so that the previous expression gets replaced by

$$\int_{c(0)=u'}^{\bar{c}(1)=e^{-i\theta} \bar{u}} D\bar{c} Dc e^{-\int_0^1 d\tau \bar{c}^\alpha \dot{c}_\alpha + \bar{c}^\alpha c_\alpha(1)} = e^{e^{-i\theta} \bar{u} \cdot u'}, \quad (3.12)$$

and (3.8) reduces to

$$\begin{aligned} \langle \phi(x, \bar{u}) \bar{\phi}(x', u') \rangle_W &= \int_0^\infty dT e^{-Tm^2} \int_0^{2\pi} \frac{d\theta}{2\pi} e^{ir\theta} \int_{x(0)=x'}^{x(1)=x} Dx \\ &\times \int_{c(0)=u'}^{\bar{c}(1)=e^{-i\theta} \bar{u}} D\bar{c} Dc \\ &\times e^{-\int_0^1 d\tau \left(\frac{1}{4T} \dot{x}^2 - ig\dot{x}^\mu W_\mu^a \bar{c}^\alpha (T^a)_\alpha^\beta c_\beta + \bar{c}^\alpha \dot{c}_\alpha \right) + \bar{c}^\alpha c_\alpha(1)}. \end{aligned} \quad (3.13)$$

Splitting as usual the fields into backgrounds and fluctuations,

$$c_\alpha(\tau) = u'_\alpha + \kappa_\alpha(\tau), \quad \kappa_\alpha(0) = 0 \quad (3.14)$$

$$\bar{c}^\alpha(\tau) = e^{-i\theta} \bar{u}^\alpha + \bar{\kappa}^\alpha(\tau), \quad \bar{\kappa}^\alpha(1) = 0, \quad (3.15)$$

the kinetic action for the fluctuations $\int_0^1 d\tau \bar{\kappa}^\alpha \dot{\kappa}_\alpha$ can be inverted to give the propagator

$$\langle \kappa_\alpha(\tau) \bar{\kappa}^\beta(\sigma) \rangle = \delta_\alpha^\beta \theta(\tau - \sigma). \quad (3.16)$$

Equation (3.13) gives the final form of our worldline representation for the scalar propagator dressed by a non-Abelian gauge field. It is gauge covariant, and we discuss the related Ward identities at the end of next section. For an arbitrary external gauge field configuration one is not able to perform the path integral exactly. On the other hand, one may use it for a suitable perturbative expansion or to get exact results for specific external field configurations.

As our main application, we specialize the external gauge potential to be given by a sum of plane waves. This allows for an exact path integral calculation, and produces a master formula for the scalar propagator with an

arbitrary number n of external (amputated) gluons lines directly attached to it, generating what we have called the ‘‘partial n -gluon scalar propagator.’’ As already discussed, the final result will not be gauge invariant, but it may be used as a starting building block to get eventually gauge invariant amplitudes. Also note that our master formula is similar to the one-loop Bern-Kosower-type of master formulas used in scalar QCD, that contain all one-particle irreducible diagrams with a scalar loop and an arbitrary number of external (amputated) gluon lines, the only difference being that instead of a loop we have an open line for the scalar particle.

Thus, in order to extract our partial n -gluon scalar propagator from the gauge-fixed worldline path integral (3.13), we write the potential as a sum of n gluons with polarizations ε_l , momenta k_l and colors a_l , i.e.

$$W_\mu(x) = \sum_{l=1}^n \varepsilon_\mu(k_l) T^{a_l} e^{ik_l \cdot x}, \quad (3.17)$$

and this allows us to read off the gluon vertex operator from the interaction term, namely

$$V_W[\varepsilon, k, a] := e^{ik \cdot x' + \varepsilon \cdot (x-x')} \int_0^1 d\tau e^{[ik \cdot (\tau(x-x') + y) + \varepsilon \cdot \dot{y}]} (e^{-i\theta} \bar{u}^\alpha + \bar{\kappa}^\alpha) (T^a)_\alpha^\beta (u_\beta + \kappa_\beta) \Big|_{\text{lin.}\varepsilon}, \quad (3.18)$$

which differs from the photon vertex operator of Eq. (2.7) in the presence of the color factor, given by the color generator T^a contracted with the auxiliary fields. The n -gluon two-scalar term reads

$$\begin{aligned} & \mathcal{A}_W(p, u; p', u'; \varepsilon_1, k_1, a_1, \dots, \varepsilon_n, k_n, a_n) \\ &= (ig)^n \int d^D x d^D x' e^{i(p \cdot x + p' \cdot x')} \times \int_0^\infty \frac{dT}{(4\pi T)^{\frac{D}{2}}} e^{-Tm^2 - \frac{1}{4T}(x-x')^2} \int_0^{2\pi} \frac{d\theta}{2\pi} e^{ir\theta + e^{-i\theta} \bar{u} \cdot u'} \prod_{l=1}^n \int_0^1 d\tau_l \\ & \times \langle V_W[k_1, \varepsilon_1, a_1] \cdots V_W[k_n, \varepsilon_n, a_n] \rangle, \end{aligned} \quad (3.19)$$

where the average is computed with the propagators (2.9) and (3.16). Due to the fact that the color part of the gluon vertex operator factors out, the integrand of the non-color part of the expression (3.19) is identical to its photon counterpart given in Ref. (2.14). We, thus, obtain the following final result for the tree-level partial n -gluon scalar propagator

$$\begin{aligned} & \tilde{\mathcal{A}}_W(p, u; p', u'; \varepsilon_1, k_1, a_1, \dots, \varepsilon_n, k_n, a_n) \\ &= (ig)^n \int_0^\infty dT e^{-T(m^2 + p^2)} \prod_{l=1}^n \int_0^1 d\tau_l \times \exp \left\{ T(p-p') \cdot \sum_l (-k_l \tau_l + i\varepsilon_l) + T \sum_{l,l'} [k_l \cdot k_{l'} \Delta_{l-l'} - 2i\varepsilon_l \cdot k_{l'} \dot{\Delta}_{l-l'} + \varepsilon_l \cdot \varepsilon_{l'} \ddot{\Delta}_{l-l'}] \right\} \Big|_{\text{m.l.}} \\ & \times \int_0^{2\pi} \frac{d\theta}{2\pi} e^{ir\theta + e^{-i\theta} \bar{u} \cdot u'} \langle (e^{-i\theta} \bar{u}^{\alpha_1} + \bar{\kappa}^{\alpha_1}(\tau_1)) (T^{a_1})_{\alpha_1}^{\beta_1} (u_{\beta_1} + \kappa_{\beta_1}(\tau_1)) \cdots \cdots (e^{-i\theta} \bar{u}^{\alpha_n} + \bar{\kappa}^{\alpha_n}(\tau_n)) (T^{a_n})_{\alpha_n}^{\beta_n} (u_{\beta_n} + \kappa_{\beta_n}(\tau_n)) \rangle, \end{aligned} \quad (3.20)$$

whose corresponding QFT Feynman diagrams have the same structure as those depicted in Fig. 1 for the n -photon amplitude. Let us underline that, since no three- and four-gluon QCD vertices are involved in our worldline computation, expression (3.20) only represents the gluon-irreducible part of the full n -gluon two-scalar amplitude, i.e. the part of the amplitude made of Feynman diagrams that cannot be parted by cutting a gluon internal line.

Before studying some special case let us consider the θ integrals present in the previous Bern-Kosower-type formula. Let us define

$$F(k, \bar{u} \cdot u') := \int_0^{2\pi} \frac{d\theta}{2\pi} e^{ik\theta + e^{-i\theta} \bar{u} \cdot u'}, \quad (3.21)$$

with $k \in \mathbb{Z}$. We can solve it by transforming it into a clockwise contour integral over the unit circle: let us define $z := e^{-i\theta}$. Hence,

$$F(k, \bar{u} \cdot u') = \oint \frac{dz}{-2\pi iz} \frac{1}{z^k} e^{\bar{u} \cdot u' z} = \begin{cases} \frac{1}{k!} (\bar{u} \cdot u')^k, & k \geq 0 \\ 0, & k < 0 \end{cases}. \quad (3.22)$$

Notice that, although the function $e^{\bar{u} \cdot u' z}$ presents an essential singularity at $z \rightarrow \infty$, it is a holomorphic function in any bounded domain, and it can be written as a power series. Thus, the contour integral over the unit circle picks out the order k coefficient of such power series.

Let us now single out a few examples in order to further clarify the results obtained.

A. Free scalar propagator

The free scalar propagator, for a field in a totally symmetric rank- r representation, can obviously be obtained from the above formalism by considering zero external gluons ($n = 0$). This will help us to fix an overall prefactor. For $n = 0$, Eq. (3.20) reduces to

$$\begin{aligned} \tilde{\mathcal{A}}_W(p, u; -p, u') &= \frac{1}{p^2 + m^2} F(r, \bar{u} \cdot u') \\ &= \frac{\bar{u}^{\alpha_1} \cdots \bar{u}^{\alpha_r} u'_{\beta_1} \cdots u'_{\beta_r} \delta_{\alpha_1 \cdots \alpha_r}^{\beta_1 \cdots \beta_r}}{r! (p^2 + m^2)}, \end{aligned} \quad (3.23)$$

where $\delta_{\alpha_1 \cdots \alpha_r}^{\beta_1 \cdots \beta_r} := \delta_{\alpha_1}^{(\beta_1} \cdots \delta_{\alpha_r}^{\beta_r)}$ is the identity in the totally symmetric rank- r representation. Equation (3.23) corresponds to the free propagator for the polarized scalar field, and one may obtain the free propagator for the unpolarized scalar field by stripping off the prefactor $\frac{\bar{u}^{\alpha_1} \cdots \bar{u}^{\alpha_r} u'_{\beta_1} \cdots u'_{\beta_r}}{r!}$ from (3.23). In fact, note that

$$\frac{\bar{u}^{\alpha_1} \cdots \bar{u}^{\alpha_r} u'_{\beta_1} \cdots u'_{\beta_r} \delta_{\alpha_1 \cdots \alpha_r}^{\beta_1 \cdots \beta_r}}{r!} = \mathbb{P}_r \langle \bar{u} | u' \rangle = \mathbb{P}_r e^{\bar{u} \cdot u'}, \quad (3.24)$$

where \mathbb{P}_r is the projector on the above irrep—basically it picks up the order r term in the Taylor expansion of the exponent. Furthermore, using the identity

$$\begin{aligned} &\frac{1}{r!} \int \prod_{\alpha=1}^N \frac{d\bar{u}^{\alpha} u_{\alpha}}{2\pi i} e^{-\bar{u} \cdot u} u_{\alpha_1} \cdots u_{\alpha_r} \bar{u}^{\beta_1} \cdots \bar{u}^{\beta_r} \\ &= \frac{1}{r!} \frac{\partial^r}{\partial \bar{v}^{\alpha_1} \cdots \partial \bar{v}^{\alpha_r}} \frac{\partial^r}{\partial v_{\beta_1} \cdots \partial v_{\beta_r}} e^{\bar{v} \cdot v} \Big|_{v=\bar{v}=0} \\ &= \delta_{\alpha_1 \cdots \alpha_r}^{\beta_1 \cdots \beta_r}, \end{aligned} \quad (3.25)$$

one can easily prove the following composition rule for the free propagator

$$\begin{aligned} &\int d^D x' \int \prod_{\alpha=1}^N \frac{d\bar{u}'^{\alpha} u'_{\alpha}}{2\pi i} e^{-\bar{u}' \cdot u'} \\ &\times \langle x, \bar{u} \left| \frac{\mathbb{P}_r}{\hat{p}^2 + m^2} \right| x', u' \rangle \langle x', \bar{u}' \left| \frac{\mathbb{P}_r}{\hat{p}^2 + m^2} \right| x'', u'' \rangle \\ &= \langle x, \bar{u} \left| \frac{\mathbb{P}_r}{(\hat{p}^2 + m^2)^2} \right| x'', u'' \rangle. \end{aligned} \quad (3.26)$$

In Eq. (3.25), we used the sources v_{α} and \bar{v}^{α} to insert the terms $u_{\alpha_1} \cdots u_{\alpha_r} \bar{u}^{\beta_1} \cdots \bar{u}^{\beta_r}$ into the Gaussian integral.

Of course the prefactor $\frac{\bar{u}^{\alpha_1} \cdots \bar{u}^{\alpha_r} u'_{\beta_1} \cdots u'_{\beta_r}}{r!}$ is universal; i.e. it only depends on the two external scalar lines and not on the gluon vertices. Therefore, the recipe to get the generic n -gluon two-scalar term from Eq. (3.20) is to strip off such color prefactors and truncate (i.e. multiply by the inverse free propagators) the external scalar lines.

B. Partial n -gluon scalar propagator

For the simplest case of one gluon, expression (3.20) reproduces the Feynman vertex linear in the gluon, whose color part is given by the expression

$$\begin{aligned} F^a(\bar{u}, u') &= \oint \frac{dz}{-2\pi iz} \frac{1}{z^r} e^{\bar{u} \cdot u' z} \\ &\times \langle (\bar{u}^{\alpha} z + \bar{\kappa}^{\alpha}(\tau)) (T^a)_{\alpha}^{\beta} (u_{\beta} + \kappa_{\beta}(\tau)) \rangle \\ &= \oint \frac{dz}{-2\pi iz} \frac{1}{z^{r-1}} e^{\bar{u} \cdot u' z} \bar{u} T^a u' \\ &= \frac{1}{(r-1)!} (\bar{u} \cdot u')^{r-1} \bar{u} T^a u' \\ &= \frac{1}{r!} \bar{u}^{\alpha_1} \cdots \bar{u}^{\alpha_r} (T^a_{(r)})^{\beta_1 \cdots \beta_r}_{\alpha_1 \cdots \alpha_r} u'_{\beta_1} \cdots u'_{\beta_r}, \end{aligned} \quad (3.27)$$

where

$$(T^a_{(r)})^{\beta_1 \cdots \beta_r}_{\alpha_1 \cdots \alpha_r} = r (T^a)_{(\alpha_1} (\beta_1 \delta_{\alpha_2}^{\beta_2} \cdots \delta_{\alpha_r}^{\beta_r}), \quad \underbrace{\square \square \square \square}_r \quad (3.28)$$

is a generator of $SU(N)$ for the rank- r totally symmetric representation. Above, we used the fact that $\langle \kappa_{\beta}(\tau) \bar{\kappa}^{\alpha}(\tau) \rangle = \frac{1}{2} \delta_{\beta}^{\alpha}$ and that T^a is traceless. Thus, it reads

$$\begin{aligned}\tilde{\mathcal{A}}_W(p, u; p', u'; \varepsilon, k, a) &= ig \int_0^\infty dT e^{-T(m^2+p^2)} \int_0^1 d\tau T e^{-T(k^2+2p \cdot k\tau)} i\varepsilon \cdot (p-p') \frac{1}{r!} \bar{u}^{\alpha_1} \cdots \bar{u}^{\alpha_r} (T_{(r)}^a)_{\alpha_1 \dots \alpha_r} \beta_1 \cdots \beta_r u'_{\beta_1} \cdots u'_{\beta_r} \\ &= ig \frac{i\varepsilon \cdot (p-p')}{(p^2+m^2)(p'^2+m^2)} (T_{(r)}^a)_{\alpha_1 \dots \alpha_r}^{\beta_1 \cdots \beta_r} \frac{\bar{u}^{\alpha_1} \cdots \bar{u}^{\alpha_r} u'_{\beta_1} \cdots u'_{\beta_r}}{r!},\end{aligned}\quad (3.29)$$

which—after truncating the external scalar lines and stripping off the factor $\frac{\bar{u} \cdots u'}{r!}$ —correctly reproduces the one-gluon two-scalar vertex

$$\tilde{\mathcal{A}}_{2s,1g}(p, \alpha; p', \beta; \varepsilon, k, a) = ig i\varepsilon \cdot (p-p') (T_{(r)}^a)_{\alpha_1 \dots \alpha_r}^{\beta_1 \cdots \beta_r}. \quad (3.30)$$

For the case of two external gluons the auxiliary field correlator involves two generators, and the color part reads

$$\begin{aligned}F^{a_1 a_2}(\bar{u}, u') &= \oint \frac{dz}{-2\pi i z} \frac{1}{z^r} e^{\bar{u} \cdot u' z} \langle ((\bar{u}^{\alpha_1} z + \bar{\kappa}^{\alpha_1}(\tau_1))(T_{(r)}^{a_1})_{\alpha_1} \beta_1 (u_{\beta_1} + \kappa_{\beta_1}(\tau_1)) (\bar{u}^{\alpha_2} z + \bar{\kappa}^{\alpha_2}(\tau_2))(T_{(r)}^{a_2})_{\alpha_2} \beta_2 (u_{\beta_2} + \kappa_{\beta_2}(\tau_2))) \\ &= \oint \frac{dz}{-2\pi i z} e^{\bar{u} \cdot u' z} \left(\frac{1}{z^{r-2}} \bar{u} T^{a_1} u' \bar{u} T^{a_2} u' + \frac{1}{z^{r-1}} \bar{u} T^{a_1} T^{a_2} u' \theta(\tau_1 - \tau_2) \right. \\ &\quad \left. + \frac{1}{z^{r-1}} \bar{u} T^{a_2} T^{a_1} u' \theta(\tau_2 - \tau_1) + \frac{1}{z} \text{tr}(T^{a_1} T^{a_2}) \theta(\tau_1 - \tau_2) \theta(\tau_2 - \tau_1) \right).\end{aligned}\quad (3.31)$$

The last term obviously vanishes for all r 's, whereas the first one is nonzero when $r \geq 2$. Furthermore, by introducing $1 = \theta(\tau_1 - \tau_2) + \theta(\tau_2 - \tau_1)$ in front of the first term, the latter reduces to

$$\begin{aligned}F^{a_1 a_2}(\bar{u}, u') &= \frac{1}{(r-1)!} [\delta_{r \geq 2}(r-1) (\bar{u} u')^{r-2} \bar{u} T^{a_1} u' \bar{u} T^{a_2} u' + (\bar{u} u')^{r-1} \bar{u} T^{a_1} T^{a_2} u'] \theta(\tau_1 - \tau_2) \\ &\quad + \frac{1}{(r-1)!} [\delta_{r \geq 2}(r-1) (\bar{u} u')^{r-2} \bar{u} T^{a_2} u' \bar{u} T^{a_1} u' + (\bar{u} u')^{r-1} \bar{u} T^{a_2} T^{a_1} u'] \theta(\tau_2 - \tau_1),\end{aligned}\quad (3.32)$$

with an obvious notation for $\delta_{r \geq 2}$. The expressions in the square brackets are just the products of the generators in the rank- r totally symmetric representation, decomposed in terms of the fundamental representation, i.e.

$$F^{a_1 a_2}(\bar{u}, u') = \frac{\bar{u}^{\alpha_1} \cdots \bar{u}^{\alpha_r} u'_{\beta_1} \cdots u'_{\beta_r}}{r!} [\theta(\tau_1 - \tau_2) (T_{(r)}^{a_1} T_{(r)}^{a_2})_{\alpha_1 \dots \alpha_r}^{\beta_1 \cdots \beta_r} + \theta(\tau_2 - \tau_1) (T_{(r)}^{a_2} T_{(r)}^{a_1})_{\alpha_1 \dots \alpha_r}^{\beta_1 \cdots \beta_r}], \quad (3.33)$$

which, once again stripping off the prefactor $\frac{\bar{u} \cdots u'}{r!}$, gives the correct color factors for the two-gluon term of the dressed scalar propagator. Finally, one can insert the latter into the formula (3.20) and get the final result for the partial two-gluon scalar propagator: the integrals over τ_1 , τ_2 and T reproduce the correct form factors. In particular the term involving the expression $\varepsilon_1 \cdot \varepsilon_2 \delta(\tau_1 - \tau_2)$ corresponds to the Feynman diagram with the two-gluon “seagull” vertex.

The partial n -gluon scalar propagator term turns out to be a straightforward generalization of the previous expressions. In particular, for an even number of gluons, the color factor displays the same features as the two-gluon one: One finds trace terms that vanish as they multiply a full product of theta functions. All the other terms combine into time-ordered products of color generators in the rank- r totally symmetric representation, i.e.

$$F^{a_1 \dots a_n}(u, u') = \frac{\bar{u}^{\alpha_1} \cdots \bar{u}^{\alpha_r} u'_{\beta_1} \cdots u'_{\beta_r}}{r!} \sum_{\sigma \in S_n} \theta(\tau_{\sigma(1)} - \tau_{\sigma(2)}) \cdots \theta(\tau_{\sigma(n-1)} - \tau_{\sigma(n)}) (T_{(r)}^{a_{\sigma(1)}} \cdots T_{(r)}^{a_{\sigma(n)}})_{\alpha_1 \dots \alpha_r}^{\beta_1 \cdots \beta_r}, \quad (3.34)$$

where S_n is the group of permutations of n elements. The same happens for an odd number of gluons.

C. Gauge-covariance of the dressed propagator and Ward identities

The worldline representation for the dressed propagator in scalar QCD, Eq. (3.13), comes in handy to show the gauge covariance of the propagator under finite transformations.

Upon a finite gauge transformation $U(x) = e^{ig\varepsilon(x)}$, with $\varepsilon(x) = \varepsilon^a(x) T^a$ and $(T^a)_\alpha^\alpha$ in the fundamental representation, the gauge field transforms as

$$\tilde{W}_\mu(x) = U(x) \left[\frac{i}{g} \partial_\mu + W_\mu(x) \right] U^\dagger(x), \quad (3.35)$$

and the worldline Lagrangian in (3.13) is gauge invariant provided the auxiliary fields transform as

$$\begin{aligned} c(\tau) &\rightarrow U(x(\tau))c(\tau) \\ \bar{c}(\tau) &\rightarrow \bar{c}(\tau)U^\dagger(x(\tau)). \end{aligned} \quad (3.36)$$

In particular, using that u' and \bar{u} are the boundary values for $c(\tau)$ and $\bar{c}(\tau)$, at $\tau = 0$ and $\tau = 1$ respectively, the previous rule implies that

$$\begin{aligned} u'_\alpha &\rightarrow (U(x'))_{\tilde{\alpha}}{}^\alpha u'_\alpha =: v'_\alpha \\ \bar{u}^\beta &\rightarrow \bar{u}^\beta (U^\dagger(x))_{\tilde{\beta}}{}^\beta =: \bar{v}^\beta, \end{aligned} \quad (3.37)$$

and

$$\langle \tilde{\phi}(x, \bar{v}) \tilde{\phi}(x', v') \rangle_{\tilde{W}} = \langle \phi(x, \bar{u}) \bar{\phi}(x', u') \rangle_W \quad (3.38)$$

for the polarized scalar propagator. Using the transformation rules (3.37) and stripping off the color factors, we obtain

$$\begin{aligned} &\langle \tilde{\phi}_{\tilde{\alpha}_1 \dots \tilde{\alpha}_r}(x) \tilde{\phi}^{\tilde{\beta}_1 \dots \tilde{\beta}_r}(x') \rangle_{\tilde{W}} \\ &= (U(x))_{\tilde{\alpha}_1}{}^{\alpha_1} \dots (U(x))_{\tilde{\alpha}_r}{}^{\alpha_r} \langle \phi_{\alpha_1 \dots \alpha_r}(x) \bar{\phi}^{\beta_1 \dots \beta_r}(x') \rangle_W \\ &\quad \times (U^\dagger(x'))_{\tilde{\beta}_1}{}^{\beta_1} \dots (U^\dagger(x'))_{\tilde{\beta}_r}{}^{\beta_r}, \end{aligned} \quad (3.39)$$

that is the correct transformation rule for the dressed propagator of a scalar field in the rank- r totally symmetric representation of $SU(N)$. At the infinitesimal level, the previous expression yields a generating functional of Ward-identities for the above amplitudes, namely

$$\begin{aligned} 0 &= D_\mu(W) \frac{\delta}{\delta W_\mu^a(y)} \langle \phi_{\alpha_1 \dots \alpha_r}(x) \bar{\phi}^{\beta_1 \dots \beta_r}(x') \rangle_W \\ &\quad + ig \delta^{(D)}(y-x) (T_{(r)}^a)_{\tilde{\alpha}_1 \dots \tilde{\alpha}_r}{}^{\tilde{\alpha}_1 \dots \tilde{\alpha}_r} \langle \phi_{\tilde{\alpha}_1 \dots \tilde{\alpha}_r}(x) \bar{\phi}^{\beta_1 \dots \beta_r}(x') \rangle_W \\ &\quad - ig \delta^{(D)}(y-x') \langle \phi_{\alpha_1 \dots \alpha_r}(x) \bar{\phi}^{\beta_1 \dots \beta_r}(x') \rangle_W (T_{(r)}^a)_{\tilde{\beta}_1 \dots \tilde{\beta}_r}{}^{\beta_1 \dots \beta_r}. \end{aligned} \quad (3.40)$$

For example, at the leading order in the coupling constant g the latter gives an identity between the one-gluon two-scalar term and the free scalar propagator. For definiteness let us consider $r = 1$. In Fourier space, and using the truncated amplitudes, the previous functional identity reduces to the perturbative Ward identity

$$\begin{aligned} 0 &= \delta^{(D)}(k+p+p') [\tilde{A}_{2s,1g}(p, \alpha; p', \beta; -ik, k, a) \\ &\quad + ig (T^a)_\alpha{}^\beta (p^2 - p'^2)], \end{aligned} \quad (3.41)$$

that does indeed vanish. Notice that in (3.41) the replacement $\varepsilon \rightarrow -ik$ implements the covariant derivative of the first term of (3.40).

IV. CONCLUSIONS AND OUTLOOK

We presented a worldline model where a scalar relativistic particle is coupled to a non-Abelian gauge field. The quantization of the model yields the scalar propagator dressed by the external gauge field. The path ordering—

needed for the gauge-covariance of the model—is realized using (commuting) auxiliary fields that are coupled to the gauge potential. The addition of a worldline Abelian gauge field along with a Chern-Simons term allows us to describe scalar particles that sit in an arbitrary rank- r symmetric representation of the gauge group. We have specialized the non-Abelian gauge field to be given by a sum of plane waves, and obtained explicit expressions for the scalar propagator dressed by an arbitrary number n of external gluons directly attached to the scalar line, the partial n -gluon scalar propagator. This is, in general, a gauge-dependent object, as three- and four-gluon vertices are not included, but it is valid off-shell and it could be used as a building block for constructing amplitudes. The resulting expressions are factorized into a color part and a kinematic part: their integral representation, provided by the present worldline model, may nicely combine with integration-by-parts techniques, that were shown to simplify the calculation of QCD form factors [43–45].

One future line of research for extending our work would be to find a simple way of adding the gluon self-interactions and obtain the complete tree-level amplitudes with 2 scalars and n gluons. One option would be to investigate a kind of “tree replacement rules” of the type used successfully in similar Bern-Kosower formulas, where the scalar particle performs a loop instead of a line.

Also, the present approach can be generalized straightforwardly to scalar QCD, with scalar particles in a rank- r totally antisymmetric representation, by using anticommuting auxiliary fields. Another quite welcome extension would be the treatment of the quark propagator in QCD, that may be obtained by coupling the external non-Abelian field to a spinning particle, i.e. by using a particle with worldline supersymmetry, where the quantization of the spinorial coordinates describe the spin degrees of freedom. An alternative approach, that needs path ordering, is the inclusion of a matrix-valued (spin-factor) potential—see e.g. Refs. [25,46,47].

A possible application of the n -gluon master formula obtained above, would be the reconstruction, in any covariant gauge, of the scalar-gluon vertex which is known as the Ball-Chiu vertex [48]. In previous studies of form factors for QED and QCD—see Refs. [31,43,45]—it was shown that the worldline formalism simplifies such calculations, and the results can be expressed in a much more compact way. Finally, it would be nice to extend the present treatment by adding the coupling to external gravity, along the lines of what has been done for the one-loop effective actions in curved space [10–12].

ACKNOWLEDGMENTS

The authors thank Idrish Huet and Christian Schubert for helpful discussions. N. A. was supported by the PRODEP (Programa para el Desarrollo Profesional Docente) Grant No. DSA/103.5/14/11184; he thanks the Università di Bologna and the Università di Modena e Reggio Emilia for their warm hospitality while parts of this work were

completed. O. C. was partly supported by the UCMEXUS–CONACYT (University of California Institute for Mexico and the United States–Consejo Nacional de Ciencia y Tecnología) Grant No. CN-12-564; he thanks the Departamento de Física of the Universidad Nacional de La Plata for its warm hospitality while parts of this work were completed.

APPENDIX: COHERENT STATE PATH INTEGRAL NORMALIZATION

Let us consider a Hilbert space spanned by a complete set of harmonic oscillator states $\{|n\rangle\}$ with $n = 0, 1, \dots, \infty$, and let \hat{c}^\dagger and \hat{c} be a pair of creation and annihilation operators acting on that Hilbert space and satisfying the canonical commutation relation $[\hat{c}, \hat{c}^\dagger] = 1$. These operators generate the harmonic oscillators states $\{|n\rangle\}$ in the usual way. Let us now define the ket and bra *coherent states* as right and left eigenstates of \hat{c} and \hat{c}^\dagger , respectively

$$\hat{c}|u\rangle = u|u\rangle, \quad \langle\bar{u}|\hat{c}^\dagger = \langle\bar{u}|\bar{u}, \quad (\text{A1})$$

where u and \bar{u} are complex numbers. They can be constructed out of the Fock vacuum as

$$|u\rangle = e^{u\hat{c}^\dagger}|0\rangle, \quad \langle\bar{u}| = \langle 0|e^{\bar{u}\hat{c}}, \quad (\text{A2})$$

and they form an overcomplete basis of the Hilbert space. Indeed their overlap reads

$$\langle\bar{u}|u'\rangle = e^{\bar{u}u'}. \quad (\text{A3})$$

After this introduction to bosonic coherent states we prove the following identity:

$$\langle\bar{u}|e^{-i\theta\hat{c}^\dagger\hat{c}}|u'\rangle = e^{\bar{u}u'e^{-i\theta}}. \quad (\text{A4})$$

A simple way of proving it is to consider the wave function $\langle\bar{u}|\psi\rangle$, corresponding to a generic state $|\psi\rangle$ of the Hilbert space. On this wave function the creation and annihilation operators act as $\hat{c}^\dagger \rightarrow \bar{u}$ and $\hat{c} \rightarrow \frac{\partial}{\partial \bar{u}}$, so that the operator $e^{-i\theta\hat{c}^\dagger\hat{c}} \rightarrow e^{-i\theta\bar{u}\frac{\partial}{\partial \bar{u}}}$ is just seen to generate a finite scaling of the \bar{u} coordinate, namely $\bar{u} \rightarrow e^{-i\theta}\bar{u}$. Then (A4) follows immediately. An alternative proof goes as follows. Let us start from the ‘‘normal order’’ rule

$$(\hat{c}^\dagger\hat{c})^m = \sum_{k=0}^m S(m, k)(\hat{c}^\dagger)^k(\hat{c})^k, \quad (\text{A5})$$

where $S(m, k)$ are the so-called ‘‘Stirling numbers of second kind’’ that are defined by¹

¹These numbers appear often in reordering problems, as for example in the worldline two-loop computation of the Euler-Heisenberg effective Lagrangian of scalar QED and spinor QED [49].

$$S(m, k) = \frac{1}{k!} \sum_{l=0}^k (-)^l \binom{k}{l} (k-l)^m, \quad (\text{A6})$$

with $S(n, 0) = \delta_{n,0}$ and $S(n, 1) = 1$. Now from the definition and properties of the coherent state and Eq. (A5), one can write the left-hand side of Eq. (A4) as

$$\begin{aligned} \langle\bar{u}|e^{-i\theta\hat{c}^\dagger\hat{c}}|u'\rangle &= \langle\bar{u}|\sum_{n=0}^{\infty} \frac{(-i\theta)^n}{n!} (a^\dagger a)^n|u'\rangle \\ &= e^{\bar{u}u'} \sum_{n=0}^{\infty} \frac{(-i\theta)^n}{n!} \sum_{k=0}^n S(n, k)(\bar{u}u')^k \\ &= e^{\bar{u}u'} \sum_{n=0}^{\infty} \frac{(-i\theta)^n}{n!} T_n(\bar{u}u'), \end{aligned} \quad (\text{A7})$$

where $T_n(x)$ are the so-called Touchard polynomials. Such polynomials can be also obtained through the exponential generating function

$$e^{x(e^z-1)} = \sum_{n=0}^{\infty} T_n(x) \frac{z^n}{n!}. \quad (\text{A8})$$

Therefore, the series in (A7) yields the right-hand side of Eq. (A4), which is thus proved.

Expression (A4) can thus be used to fix the normalization for the harmonic oscillator coherent state path integral. A free coherent state path integral can be built up by inserting spectral decompositions of the identity

$$\mathbb{1} = \int \frac{d\bar{c}dc}{2\pi i} e^{-\bar{c}c|c\rangle\langle\bar{c}|} \quad (\text{A9})$$

into the expression (A3) for the scalar product. One gets

$$\begin{aligned} \langle\bar{u}|u'\rangle &= e^{\bar{u}u'} \\ &= \int \prod_{k=1}^{n-1} \frac{d\bar{c}_k dc_k}{2\pi i} \exp \left\{ \bar{c}_n c_{n-1} - \sum_{k=1}^{n-1} \bar{c}_k (c_k - c_{k-1}) \right\}, \end{aligned} \quad (\text{A10})$$

where we defined $c_0 := u'$ and $\bar{c}_n := \bar{u}$. Furthermore, we define a time parameter $0 \leq \tau \leq 1$ and identify $c(\tau_k) := c_k$ and $\bar{c}(\tau_k) := \bar{c}_k$, with $\tau_k - \tau_{k-1} = \frac{1}{n-1} =: \epsilon$. Thus, in the large n limit, we may identify the latter as the free coherent state path integral

$$\langle\bar{u}|u'\rangle = e^{\bar{u}u'} = \int_{c(0)=u'}^{\bar{c}(1)=\bar{u}} D\bar{c}Dc e^{-S_f[c, \bar{c}]}, \quad (\text{A11})$$

with

$$S_f[c, \bar{c}] = \int_0^1 d\tau \bar{c}(\tau) \dot{c}(\tau) - \bar{c}c(1). \quad (\text{A12})$$

We may, thus, split the paths into backgrounds, satisfying the free equations of motion $\dot{\bar{c}} = \dot{c} = 0$ with corresponding boundary conditions, and quantum fluctuations

$$c(\tau) = u' + \kappa(\tau), \quad \kappa(0) = 0 \quad (\text{A13})$$

$$\bar{c}(\tau) = \bar{u} + \bar{\kappa}(\tau), \quad \bar{\kappa}(1) = 0. \quad (\text{A14})$$

Hence,

$$\langle \bar{u} | u' \rangle = e^{\bar{u}u'} = e^{\bar{u}u'} \int_{\kappa(0)=0}^{\bar{\kappa}(1)=0} D\bar{\kappa}D\kappa e^{-\int_0^1 d\tau \bar{\kappa} \dot{\kappa}}, \quad (\text{A15})$$

which in turn gives

$$\int_{\kappa(0)=0}^{\bar{\kappa}(1)=0} D\bar{\kappa}D\kappa e^{-\int_0^1 d\tau \bar{\kappa} \dot{\kappa}} = 1. \quad (\text{A16})$$

Let us now consider the path integral

$$Z(\bar{u}, u'; \theta) := \int_{c(0)=u'}^{\bar{c}(1)=\bar{u}} D\bar{c}Dc e^{-S[c, \bar{c}; \theta]}, \quad (\text{A17})$$

with

$$S[c, \bar{c}] = \int_0^1 d\tau \bar{c}(\tau)(\partial_\tau + i\theta)c(\tau) - \bar{c}c(1). \quad (\text{A18})$$

We can solve the latter in a similar way as above, i.e. by splitting the paths into backgrounds,

$$(\partial_\tau + i\theta)c(\tau) = 0, \quad c(0) = u' \Rightarrow C(\tau) = u' e^{-i\theta\tau} \quad (\text{A19})$$

$$(-\partial_\tau + i\theta)\bar{c}(\tau) = 0, \quad \bar{c}(1) = \bar{u} \Rightarrow \bar{C}(\tau) = \bar{u} e^{i\theta(\tau-1)}, \quad (\text{A20})$$

and quantum fluctuations $\kappa(\tau)$ and $\bar{\kappa}(\tau)$ with the same boundary conditions as above. By setting

$$c(\tau) = e^{-i\theta\tau}(u' + \kappa(\tau)) \quad (\text{A21})$$

$$\bar{c}(\tau) = e^{i\theta\tau}(e^{-i\theta}\bar{u} + \bar{\kappa}(\tau)), \quad (\text{A22})$$

one gets

$$\begin{aligned} Z(\bar{u}, u'; \theta) &:= e^{e^{-i\theta}\bar{u}u'} \int_{\kappa(0)=0}^{\bar{\kappa}(1)=\bar{0}} D\bar{\kappa}D\kappa e^{-\int_0^1 d\tau \bar{\kappa} \dot{\kappa}} \\ &= e^{e^{-i\theta}\bar{u}u'}, \end{aligned} \quad (\text{A23})$$

where we made use of (A16). In summary,

$$\int_{c(0)=u'}^{\bar{c}(1)=\bar{u}} D\bar{c}Dc e^{-\int_0^1 d\tau \bar{c}(\tau)(\partial_\tau + i\theta)c(\tau) + \bar{c}c(1)} = e^{e^{-i\theta}\bar{u}u'}, \quad (\text{A24})$$

which we thus identify with (A4). Thus, by adding a constant term to the action and considering $\alpha = 1, \dots, N$ independent pairs of oscillator operators, we finally get

$$\begin{aligned} \int_{c(0)=u'}^{\bar{c}(1)=\bar{u}} D\bar{c}Dc e^{-\int_0^1 d\tau \bar{c}^\alpha(\tau)(\partial_\tau + i\theta)c_\alpha(\tau) + \bar{c} \cdot c(1) + i\theta r} \\ = \langle \bar{u} | e^{-i\theta(\hat{c}^\dagger \cdot \hat{c} - r)} | u' \rangle = e^{e^{-i\theta}\bar{u} \cdot u' + i\theta r}, \end{aligned} \quad (\text{A25})$$

where $\mathbb{N} := \hat{c}^\dagger \cdot \hat{c} = \hat{c}^\dagger \alpha \hat{c}_\alpha$ is the total occupation number operator for the system of N harmonic oscillators.

If, on the other hand, one were to identify

$$\begin{aligned} \int_{c(0)=u'}^{\bar{c}(1)=\bar{u}} D\bar{c}Dc e^{-\int_0^1 d\tau \bar{c}^\alpha(\tau)(\partial_\tau + i\theta)c_\alpha(\tau) + \bar{c} \cdot c(1)} \\ = \langle \bar{u} | e^{-i\theta(\hat{c}^\dagger \cdot \hat{c})_s} | u' \rangle, \end{aligned} \quad (\text{A26})$$

with

$$(\hat{c}^\dagger \cdot \hat{c})_s := \frac{1}{2}(\hat{c}^\dagger \cdot \hat{c} + \hat{c} \cdot \hat{c}^\dagger) = \hat{c}^\dagger \cdot \hat{c} + \frac{N}{2} \quad (\text{A27})$$

being the symmetrized product, one would then get

$$\int_{c(0)=u'}^{\bar{c}(1)=\bar{u}} D\bar{c}Dc e^{-\int_0^1 d\tau \bar{c}^\alpha(\tau)(\partial_\tau + i\theta)c_\alpha(\tau) + \bar{c} \cdot c(1)} = e^{e^{-i\theta}\bar{u} \cdot u' - i\theta \frac{N}{2}}, \quad (\text{A28})$$

and

$$\begin{aligned} \int_{c(0)=u'}^{\bar{c}(1)=\bar{u}} D\bar{c}Dc e^{-\int_0^1 d\tau \bar{c}^\alpha(\tau)(\partial_\tau + i\theta)c_\alpha(\tau) + \bar{c} \cdot c(1) + i\theta s} \\ = \langle \bar{u} | e^{-i\theta((\hat{c}^\dagger \cdot \hat{c})_s - s)} | u' \rangle = e^{e^{-i\theta}\bar{u} \cdot u' + i\theta(s - \frac{N}{2})}. \end{aligned} \quad (\text{A29})$$

However, setting the eigenvalues of the occupation numbers to be equal,

$$(\hat{c}^\dagger \cdot \hat{c})_s - s = \hat{c}^\dagger \cdot \hat{c} - r \Rightarrow s = r + \frac{N}{2}, \quad (\text{A30})$$

the right-hand sides of (A25) and (A29) coincide.

- [1] M. J. Strassler, Field theory without Feynman diagrams: One loop effective actions, *Nucl. Phys.* **B385**, 145 (1992).
- [2] Z. Bern and D. A. Kosower, Efficient Calculation of One Loop QCD Amplitudes, *Phys. Rev. Lett.* **66**, 1669 (1991).
- [3] Z. Bern and D. A. Kosower, The computation of loop amplitudes in gauge theories, *Nucl. Phys.* **B379**, 451 (1992).
- [4] C. Schubert, Perturbative quantum field theory in the string inspired formalism, *Phys. Rep.* **355**, 73 (2001).
- [5] M. G. Schmidt and C. Schubert, Multiloop calculations in the string inspired formalism: the single spinor loop in QED, *Phys. Rev. D* **53**, 2150 (1996).
- [6] M. Reuter, M. G. Schmidt, and C. Schubert, Constant external fields in gauge theory and the spin 0, 1/2, 1 path integrals, *Ann. Phys. (N.Y.)* **259**, 313 (1997).
- [7] H. Gies, J. Sanchez-Guillen, and R. A. Vazquez, Quantum effective actions from nonperturbative worldline dynamics, *J. High Energy Phys.* **08** (2005) 067.
- [8] G. V. Dunne and C. Schubert, Worldline instantons and pair production in inhomogeneous fields, *Phys. Rev. D* **72**, 105004 (2005).
- [9] H. Gies, K. Langfeld, and L. Moyaerts, Casimir effect on the worldline, *J. High Energy Phys.* **06** (2003) 018.
- [10] F. Bastianelli and A. Zirotti, Worldline formalism in a gravitational background, *Nucl. Phys.* **B642**, 372 (2002).
- [11] F. Bastianelli, O. Corradini, and A. Zirotti, Dimensional regularization for $N = 1$ supersymmetric sigma models and the worldline formalism, *Phys. Rev. D* **67**, 104009 (2003).
- [12] F. Bastianelli, P. Benincasa, and S. Giombi, Worldline approach to vector and antisymmetric tensor fields, *J. High Energy Phys.* **04** (2005) 010.
- [13] F. Bastianelli and C. Schubert, One loop photon-graviton mixing in an electromagnetic field: Part 1, *J. High Energy Phys.* **02** (2005) 069.
- [14] F. Bastianelli, O. Corradini, and E. Latini, Higher spin fields from a worldline perspective, *J. High Energy Phys.* **02** (2007) 072.
- [15] F. Bastianelli, O. Corradini, and E. Latini, Spinning particles and higher spin fields on (A)dS backgrounds, *J. High Energy Phys.* **11** (2008) 054.
- [16] F. Bastianelli, R. Bonezzi, O. Corradini, and E. Latini, Effective action for higher spin fields on (A)dS backgrounds, *J. High Energy Phys.* **12** (2012) 113.
- [17] P. Dai, Y.-t. Huang, and W. Siegel, Worldgraph approach to Yang-Mills amplitudes from $N = 2$ spinning particle, *J. High Energy Phys.* **10** (2008) 027.
- [18] D. Fliegner, M. G. Schmidt, and C. Schubert, The higher derivative expansion of the effective action by the string inspired method. Part 1., *Z. Phys. C* **64**, 111 (1994).
- [19] D. Fliegner, P. Haberl, M. G. Schmidt, and C. Schubert, The higher derivative expansion of the effective action by the string inspired method. Part 2, *Ann. Phys. (N.Y.)* **264**, 51 (1998).
- [20] F. Bastianelli, O. Corradini, P. A. G. Pisani, and C. Schubert, Scalar heat kernel with boundary in the worldline formalism, *J. High Energy Phys.* **10** (2008) 095.
- [21] Y. Kiem, S. J. Rey, H. T. Sato, and J. T. Yee, Anatomy of one loop effective action in noncommutative scalar field theories, *Eur. Phys. J. C* **22**, 757 (2002).
- [22] R. Bonezzi, O. Corradini, S. A. Franchino Vinas, and P. A. G. Pisani, Worldline approach to noncommutative field theory, *J. Phys. A* **45**, 405401 (2012).
- [23] P. Mansfield, The fermion content of the Standard Model from a simple world-line theory, *Phys. Lett. B* **743**, 353 (2015).
- [24] J. P. Edwards, Unified theory in the worldline approach, *Phys. Lett. B* **750**, 312 (2015).
- [25] R. P. Feynman, An operator calculus having applications in quantum electrodynamics, *Phys. Rev.* **84**, 108 (1951).
- [26] K. Daikouji, M. Shino, and Y. Sumino, Bern-Kosower rule for scalar QED, *Phys. Rev. D* **53**, 4598 (1996).
- [27] F. Bastianelli, A. Huet, C. Schubert, R. Thakur, and A. Weber, Integral representations combining ladders and crossed-ladders, *J. High Energy Phys.* **07** (2014) 066.
- [28] Z. Bern, J. J. M. Carrasco, and H. Johansson, New relations for gauge-theory amplitudes, *Phys. Rev. D* **78**, 085011 (2008).
- [29] E. S. Fradkin, Concerning some general relations of quantum electrodynamics, *Zh. Eksp. Teor. Fiz.* **29**, 258 (1955) [*Sov. Phys. JETP* **2**, 361 (1956)].
- [30] L. D. Landau and I. M. Khalatnikov, The gauge transformation of the Green function for charged particles, *Zh. Eksp. Teor. Fiz.* **29**, 89 (1955) [*Sov. Phys. JETP* **2**, 69 (1956)].
- [31] N. Ahmadiniaz, A. Bashir, and C. Schubert, Multiphoton amplitudes and generalized LKF transformation in Scalar QED, [arXiv:1511.05087](https://arxiv.org/abs/1511.05087).
- [32] S. D. Badger, E. W. N. Glover, V. V. Khoze, and P. Svrcek, Recursion relations for gauge theory amplitudes with massive particles, *J. High Energy Phys.* **07** (2005) 025.
- [33] D. Forde and D. A. Kosower, All-multiplicity amplitudes with massive scalars, *Phys. Rev. D* **73**, 065007 (2006).
- [34] P. Ferrario, G. Rodrigo, and P. Talavera, Compact Multi-gluonic Scattering Amplitudes with Heavy Scalars and Fermions, *Phys. Rev. Lett.* **96**, 182001 (2006).
- [35] R. Boels and C. Schwinn, CSW rules for a massive scalar, *Phys. Lett. B* **662**, 80 (2008).
- [36] F. Bastianelli, R. Bonezzi, O. Corradini, and E. Latini, Particles with non-Abelian charges, *J. High Energy Phys.* **10** (2013) 098.
- [37] F. Bastianelli, R. Bonezzi, O. Corradini, E. Latini, and K. H. Ould-Lahoucine, A worldline approach to colored particles, [arXiv:1504.03617](https://arxiv.org/abs/1504.03617).
- [38] A. P. Balachandran, P. Salomonson, B. S. Skagerstam, and J. O. Winnberg, Classical description of particle interacting with nonabelian gauge field, *Phys. Rev. D* **15**, 2308 (1977).
- [39] A. Barducci, R. Casalbuoni, and L. Lusanna, Classical scalar and spinning particles interacting with external Yang-Mills fields, *Nucl. Phys.* **B124**, 93 (1977).
- [40] E. D'Hoker and D. G. Gagne, Worldline path integrals for fermions with general couplings, *Nucl. Phys.* **B467**, 297 (1996).
- [41] E. Fradkin, Application of functional methods in quantum field theory and quantum statistics (II), *Nucl. Phys.* **76**, 588 (1966).
- [42] P. S. Howe, S. Penati, M. Pernici, and P. K. Townsend, Wave equations for arbitrary spin from quantization of the

- extended supersymmetric spinning particle, *Phys. Lett. B* **215**, 555 (1988).
- [43] N. Ahmadinia and C. Schubert, A covariant representation of the Ball-Chiu vertex, *Nucl. Phys.* **B869**, 417 (2013).
- [44] N. Ahmadinia, C. Schubert, and V. M. Villanueva, String-inspired representations of photon/gluon amplitudes, *J. High Energy Phys.* 01 (2013) 132.
- [45] N. Ahmadinia and C. Schubert, Form factor decomposition of the off-shell four-gluon amplitudes, *Proc. Sci. QCD*, -TNT-III (2013) 002.
- [46] F. Bastianelli and P. van Nieuwenhuizen, Trace anomalies from quantum mechanics, *Nucl. Phys.* **B389**, 53 (1993).
- [47] D. M. Gitman and S. I. Zlatev, Spin factor in path integral representation for Dirac propagator in external fields, *Phys. Rev. D* **55**, 7701 (1997).
- [48] J. S. Ball and T-W. Chiu, Analytic properties of the vertex function in gauge theories. II, *Phys. Rev. D* **22**, 2550 (1980); **23**, 3085(E) (1981).
- [49] G. V. Dunne and C. Schubert, Two loop selfdual Euler-Heisenberg Lagrangians. 2. Imaginary part and Borel analysis, *J. High Energy Phys.* 06 (2002) 042.