

REVIEW

The short-lived African turquoise killifish: an emerging experimental model for ageing

Yumi Kim^{1,2}, Hong Gil Nam^{2,3} and Dario Riccardo Valenzano^{1,*}

ABSTRACT

Human ageing is a fundamental biological process that leads to functional decay, increased risk for various diseases and, ultimately, death. Some of the basic biological mechanisms underlying human ageing are shared with other organisms; thus, animal models have been invaluable in providing key mechanistic and molecular insights into the common bases of biological ageing. In this Review, we briefly summarise the major applications of the most commonly used model organisms adopted in ageing research and highlight their relevance in understanding human ageing. We compare the strengths and limitations of different model organisms and discuss in detail an emerging ageing model, the short-lived African turquoise killifish. We review the recent progress made in using the turquoise killifish to study the biology of ageing and discuss potential future applications of this promising animal model.

KEY WORDS: Ageing, Longevity, Age-associated diseases, Model organisms, Turquoise killifish, *Nothobranchius furzeri*

Introduction

Biological ageing consists of a wide range of dynamic changes that occur throughout an organism's lifespan that negatively impact all fundamental biological processes and eventually result in the loss of organismal homeostasis and, ultimately, lead to death (Kirkwood and Austad, 2000; Lopez-Otin et al., 2013). Human ageing is associated with characteristic macroscopic changes, which include hair greying, wrinkling of the skin, muscle loss and physical weakness. As individuals age they become more susceptible to a wide range of diseases. In particular, heart disease, cancer, stroke, chronic lower respiratory disease, type 2 diabetes and neurodegeneration are the most common age-associated diseases, and each represents a leading cause of death in aged individuals (Akushevich et al., 2013; Brody and Grant, 2001; Craig et al., 2015; WHO, 2011). Age-associated phenotypes are thought to result from the progressive accumulation of molecular damage, and this phenomenon is postulated to be the consequence of the age-dependent decrease in the force of selection, which fails to remove deleterious mutations that affect aspects of later life (post-fertility) (Bailey et al., 2004; Charlesworth, 2000; Dolle et al., 2000). The age-dependent accumulation of molecular damage induces decreased DNA or protein stability, failure in energy production and utilization, and disruption of homeostasis, leading to structural

and functional decay (Lopez-Otin et al., 2013). It is also predicted that mutations providing an overall fitness benefit throughout an organism's lifespan are likely to increase in frequency in a population, even if their phenotypic effect at older ages is detrimental (Williams, 1957).

Human progeroid syndromes and extreme human longevity (see definitions below) offer two biological extremes that have helped to shed light on the basic genetic and physiological mechanisms associated with accelerated ageing and extreme lifespan in humans (Burtner and Kennedy, 2010; Eisenstein, 2012). Human progeroid syndromes are a set of monogenic disorders associated with dysfunctions in the DNA repair machinery or improper formation of the nuclear lamina that lead to premature ageing-like symptoms (Burtner and Kennedy, 2010; De Sandre-Giovannoli et al., 2003; Kitano, 2014; Veith and Mangerich, 2015). Human extreme longevity is a complex phenotype that depends on the interaction between multiple genetic variants and environmental conditions. Importantly, the causal role of different biological mechanisms and genetic variants on human longevity remains elusive owing to obvious experimental limitations with human subjects and the low frequency of centenarians.

Human cell lines provide a great resource for ageing research because they allow the study of several aspects of human cellular biology in a Petri dish (de Magalhaes, 2004; Hashizume et al., 2015); however, they restrict the relevance of the findings to the cellular aspects of individual ageing, and provide limited contribution to the understanding of the *in vivo* mechanisms involved in organismal ageing. An obvious alternative strategy to overcome some of these limitations involves the use of model organisms that either share or mimic the ageing-associated processes of humans.

The use of model organisms has been key to improving the understanding of the molecular mechanisms underlying ageing and the wide spectrum of age-related diseases. The most successful model organisms used in the field of ageing include non-vertebrate models, e.g. yeast, worms and flies, and vertebrate models, e.g. zebrafish and mice. The lifespan of these model organisms in captivity ranges from a few weeks to several years, and the spectrum of their ageing phenotypes is extremely diverse, each mimicking different features associated with human ageing (Table 1). In this article, we first introduce the most commonly adopted model organisms in research on ageing and age-associated diseases. We then detail features of the African turquoise killifish that make it a promising complementary model system for the exploration of ageing *in vivo*, and highlight its potential to provide insight into ageing-related diseases.

Common model organisms used in ageing research

Non-vertebrate models

The budding yeast (*Saccharomyces cerevisiae*) is a unicellular eukaryote that is widely used in ageing research (Henderson and

¹Max Planck Institute for Biology of Ageing, D50931, Cologne, Germany.

²Department of New Biology, DGIST, 711-873, Daegu, Republic of Korea. ³Center for Plant Aging Research, Institute for Basic Science, 711-873, Daegu, Republic of Korea.

*Author for correspondence (Dario.Valenzano@age.mpg.de)

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

Table 1. Model systems used to study ageing

Model organism	Typical lifespan duration	Characteristic ageing phenotypes	Genetic interventions	No. ageing mutants to date	References
Yeast	5-14 days	<ul style="list-style-type: none"> • Nucleolus fragmentation • Failure of sporulation • Accumulation of extrachromosomal rDNA circles • Accumulation of DNA mutation • Increased apoptosis • Mitochondrial dysfunction • Reduced vacuolar acidity • Changes in protein expression and localisation • Age-dependent gene expression 	<ul style="list-style-type: none"> • Homologous recombination • Chemical mutagenesis (EMS, MS, NNG, NA, ICR-170) • Radiation (UV, X-ray) • RNAi • Targeted genome editing (CRISPR/Cas9 system, TALEN) 	825	Burhans and Weinberger, 2012; Cagney, 2009; Craig et al., 2015; Herker et al., 2004; Klass, 1977; Lewinska et al., 2014; Longo et al., 1997; Longo et al., 1996; Sinclair and Guarente, 1997; Tacutu et al., 2013; Wei et al., 2009
Worm	12-18 days at 20°C	<ul style="list-style-type: none"> • Decrease in organisation of pharynx/head • Bacterial accumulation in pharynx/intestine • Decrease in organisation and loss of sarcomeric density of body wall muscles • Decreased organisation of germ line • Decreased pharyngeal pumping • Progressive decrease of body movement • Decreased maximum velocity • Decreased response to chemotaxis • Decreased isothermal tracking • Decreased rate of defecation • Decreased progeny production • Lipofuscin accumulation • Accumulation of DNA damage • Increase in levels of carbonyl contents • Accumulation of yolk protein in body cavity • Decreased metabolic activity • Decrease in protein tyrosine kinase activity • Increase in lysosomal hydrolases • Changes in gene expression 	<ul style="list-style-type: none"> • Insertional mutagenesis (Mos1, Tc1, MosDEL, MosTIC) • Random chemical mutation (EMS, DES, ENU, formaldehyde) • Radiation (UV, X-ray) • RNAi • Targeted genome editing (CRISPR/Cas9 system, TALEN, ZFN) 	741	Collins et al., 2008; Craig et al., 2015; Croll et al., 1977; Girard et al., 2007; Hahm et al., 2015; Hamilton et al., 2005; Hertweck et al., 2003; Johnson, 2003; Johnson and Hutchinson, 1993; Klass, 1977; Kutscher and Shaham, 2014; Tacutu et al., 2013
Fly	30-40 days	<ul style="list-style-type: none"> • Decreased body mass and thorax mass • Decreased locomotor activity • Decreased climbing and flying activity • Decreased wingbeat frequency • Reduced circadian activity • Decreased reproductive capacity • Accumulation of DNA damage • Decreased metabolic rate • Mitochondrial dysfunction • Lipofuscin accumulation • Peroxide accumulation 	<ul style="list-style-type: none"> • Insertional mutagenesis (P element, piggyBac, Minos, Tol2) • Random chemical mutation (EMS, ENU, DHPLC, HMPA) • Homologous recombination • Morpholino • RNAi • Targeted genome editing (CRISPR/Cas9 system, TALEN, ZFN) 	140	Armstrong et al., 1978; Craig et al., 2015; Curtsinger et al., 1992; Ferguson et al., 2005; Grotewiel et al., 2005; Helfand and Rogina, 2003; Jacobson et al., 2010; Lamb, 1968; Lane et al., 2014; Promislow et al., 1996; Tacutu et al., 2013; Tatar et al., 1996; Walter et al., 2007
Zebrafish	36-42 months	<ul style="list-style-type: none"> • Spinal curvature • Decreased regenerative capacity • Muscle degeneration • Age-associated alterations in circadian rhythmicity 	<ul style="list-style-type: none"> • Insertional mutagenesis (Tol2, SB) • Random chemical mutation (ENU) 	19	Almaida-Pagan et al., 2014; Anchin et al., 2011; Gerhard et al., 2002; Gilbert et al., 2014; Kishi et al., 2008; Kishi et al., 2003; Lawson and Wolfe, 2011;

Continued

Table 1. Continued

Model organism	Typical lifespan duration	Characteristic ageing phenotypes	Genetic interventions	No. ageing mutants to date	References
		<ul style="list-style-type: none"> • Decreased reproductive capacity • Decreased swimming performance • Increased SA-β-galactosidase activity • Lipofuscin accumulation • Accumulation of oxidised proteins in muscle • Drusen-like lesions in retinal pigment epithelium • Telomere shortening • Changes in mitochondrial membrane composition • Decreased cytosine methylation in somatic cells 	<ul style="list-style-type: none"> • RNAi • Morpholino • Targeted genome editing (CRISPR/Cas9 system, TALEN, ZFN) 		Moens et al., 2008; Ruzicka et al., 2015
Killifish	9-26 weeks	<ul style="list-style-type: none"> • Spinal curvature • Colour loss • Emaciation • Reduced locomotor activity • Cognitive impairment • Decreased fecundity • Accumulation of lipofuscin • Increased apoptosis • Liver neoplasias • Telomere shortening • Mitochondrial impairment • Neurodegeneration • Decreased fin regeneration • Age-dependent gene expression 	<ul style="list-style-type: none"> • Insertional mutagenesis (Tol2) • Targeted genome editing (CRISPR/Cas9 system) 	6	Baumgart et al., 2015; Hartmann and Englert, 2012; Hartmann et al., 2009, 2011; Terzibasi et al., 2008; Tozzini et al., 2012; Valdesalici and Cellierino, 2003; Valenzano et al., 2011; Wendler et al., 2015
Mouse	2-3 years	<ul style="list-style-type: none"> • Body weight declines • Bone loss • Growth plate closure • Skin ulcerations • Skin atrophy • Hair greying • Alopecia • Decreased wound healing • Retinal degeneration • Ability to acquire motor skills decreases • Balance decreases • Grip strength decreases • Decreased exploratory behaviour • Impaired memory consolidation • No clear changes in anxiety but decrease in ambulation • Loss of subcutaneous adipose skin layer • Increased cancer incidence • Cellular senescence • Increased sensitivity to carcinogens • Accumulation of DNA damage • Neurodegeneration • Neural stem cell pool declines • Osteoblast differentiation decreases • Pancreatic β-cell replication decreases • Telomere shortening 	<ul style="list-style-type: none"> • Homologous recombination • Insertional mutagenesis (Tol2) • RNAi • Morpholino • Targeted genome editing (CRISPR/Cas9 system, TALEN, ZFN) 	112	Craig et al., 2015; Demetrius, 2006; Echigoya et al., 2015; Fahlstrom et al., 2011; Hasty et al., 2003; Ingram, 1983; Johnson et al., 2005; Sung et al., 2013; Tacutu et al., 2013; Vanhooren and Libert, 2013; Wang et al., 2013; Yang et al., 2014; Yuan et al., 2011, 2009; Zahn et al., 2007

CRISPR, clustered regularly-interspaced short palindromic repeats; DES, diethyl sulfate; DHPLC, denaturing high pressure liquid chromatography; EMS, ethyl methanesulfonate; ENU, N-ethyl-N-nitrosourea; HMPA, hexamethylphosphoramide; ICR-170, 2-methoxy-6-chloro-9-[3-(ethyl-2-chloroethyl)aminopropylamino] acridine dihydrochloride; MosDel, Mos1-mediated deletion; MosTIC, Mos1-excision-induced transgene-instructed gene conversion; MS, methyl methanesulfonate; NA, nitrous acid; NNG, nitrosoguanidine; RNAi, RNA interference; SA, senescence-associated; SB, sleeping beauty; TALEN, transcription activator-like effector nuclease; UV, ultraviolet; ZFN, zinc-finger nuclease.

Gottschling, 2008; Herker et al., 2004; Sinclair and Guarente, 1997; Verduyck et al., 2016). The budding yeast undergoes a limited number of replication events throughout its lifetime. The total number of daughter cells produced by a mother cell defines the yeast cell replicative lifespan (RLS), which is used as a direct measure of yeast survival. Another factor used to measure yeast survival is the survival time of populations of non-dividing yeast cells, which defines the so-called chronological life span (CLS) (Herker et al., 2004; Kaerberlein, 2010; Longo et al., 1996).

Dietary restriction (DR), i.e. reduced food intake without malnutrition (Colman et al., 2014; Piper et al., 2011), is one of the better-studied interventions capable of delaying ageing and prolonging lifespan in several species (Fontana and Partridge, 2015). In line with this, DR has been shown to increase yeast RLS as well as CLS. Some of the most important cellular effectors involved in increasing lifespan belong to the target of rapamycin (TOR) molecular pathway (Bonawitz et al., 2007; Kaerberlein et al., 2005; Powers et al., 2006). This molecular pathway is involved in the regulation of protein synthesis and degradation in response to nutrient quantity and quality (Stanfel et al., 2009), and is largely conserved from yeast to mammals. Importantly, this molecular pathway is impaired in human metabolic dysfunctions such as diabetes and obesity (Cota et al., 2008; Fraenkel et al., 2008).

Given the conservation of the basic eukaryotic intracellular organelles and machinery, yeast has been successfully used as a model to study the intracellular effects of mutated human genes involved in several age-associated neurodegenerative diseases, including Parkinson's and Alzheimer's diseases (Cooper et al., 2006; Khurana and Lindquist, 2010). However, because yeast are unicellular organisms, their use as a model system for ageing is limited to the understanding of the cellular mechanisms. Additionally, some features associated with yeast ageing are specific to yeast biology, such as the age-related accumulation of extrachromosomal ribosomal DNA (rDNA) circles (ERCs) in yeast mother cells (Sinclair and Guarente, 1997). ERCs do not have a direct correlate with the ageing process of other organisms, and demonstrate that, although some ageing-related mechanisms are shared across different taxa, some others are species-specific.

Caenorhabditis elegans (*C. elegans*) is a transparent soil nematode that has been utilized as an experimental model system since 1974 (Brenner, 1974). Its usefulness as a model can be in part attributed to the fact that it is a multicellular organism and its life cycle can be entirely recapitulated in a Petri dish. In laboratory conditions, this worm, of about 1 mm in length in the adult form, lives only a few weeks and its life cycle has been thoroughly investigated (<http://www.wormatlas.org/>). Many ageing phenotypes of the worm are also shared with other organisms, including humans, such as decreased overall body motility and food consumption (measured as pharyngeal pumping rate), progressively increased DNA damage, decreased metabolic activity, accumulation of age-pigments and dramatic changes in age-dependent gene expression (Collins et al., 2008). Interestingly, during development, *C. elegans* can enter a stress-resistant biological state called dauer, characterised by typical morphological changes and the capacity to survive through starvation, temperature changes and other stressors (Gottlieb and Ruvkun, 1994; Lithgow et al., 1995; Riddle et al., 1981). The genes involved in the regulation of dauer formation in *C. elegans* play a key role in regulating worm longevity, and their function in regulating stress responses is conserved across many organisms, including humans (Kenyon et al., 1993). The worm is highly amenable to genetic manipulation (Klass, 1983), which has enabled several genetic screens that have brought to light pathways involved

in ageing and lifespan. Large-scale screening of several worm mutant lines and genes (Klass, 1983; Wong et al., 1995; Yang and Wilson, 2000), particularly using RNA interference (RNAi) screening (Hamilton et al., 2005; Lee et al., 2003; Yanos et al., 2012), has been a key contributor to the fundamental insights made into the molecular genetics of ageing. Importantly, these genetic screens provided the first evidence that a single gene can modulate longevity in a multicellular eukaryote (Kimura et al., 1997; Klass, 1983; Lee et al., 2001; Ogg et al., 1997; Tissenbaum and Ruvkun, 1998). These results laid the foundation for the discovery of shared cellular and organismal mechanisms, such as the stress response and nutrient-sensing, that control longevity in multiple organisms, and these findings are likely to be relevant to human ageing (Kenyon, 2010; Kim, 2007; Lopez-Otin et al., 2013).

The fruit fly, *Drosophila melanogaster* (*D. melanogaster*), is a well-established model organism, with many key attributes that make it a valuable experimental system, such as being easy to maintain and amenable to manipulation using advanced genetic tools. In addition, the fruit fly community benefits from the availability of public resources, including mutant and gene libraries (<http://flybase.org/>) (Matthews et al., 2005; Millburn et al., 2016). Wild-type *D. melanogaster* can survive for just a few months in captivity, and it is therefore an excellent experimental model in which to study the biology of ageing and the effects of different interventions on overall life expectancy. Although modern flies and worms are phylogenetically equally related to vertebrates (Masoro and Austad, 2006), flies have some features that make them more closely resemble higher vertebrates, such as a complex and centralised brain, a heart (Chintapalli et al., 2007), and the presence of multipotent adult stem cells in the mid-gut and the gonads (Micchelli and Perrimon, 2006; Ohlstein and Spradling, 2006; Wallenfang et al., 2006). Their functional proximity to vertebrates has allowed the development of several fly models that are useful for the study of mammalian ageing, in particular the effects of ageing on muscle, brain, cardiac and intestinal tissues (Demontis and Perrimon, 2010; Guo et al., 2014; Haddadi et al., 2014; Ocorr et al., 2007).

The availability of rapid and effective molecular, biochemical and genetic tools for application in invertebrate models has provided a great advantage in the field of ageing. Indeed, research using these models has yielded important insights into the basic cellular and molecular mechanisms that underlie the ageing process in a wide range of living organisms. However, invertebrate model organisms do not allow exploration of all aspects of human ageing. For instance, the currently available invertebrate model organisms (i.e. worms and flies) are not ideal platforms to study the process of tumorigenesis and cancer during ageing, because adults of these organisms are mostly composed of post-mitotic cells, i.e. cells that no longer undergo the cell cycle and therefore cannot develop cancer (Masoro, 1996). Additionally, although they are equipped with an effective innate immune system, invertebrates lack a vertebrate-like lymphocyte-based adaptive immune system (Johnson, 2003). In addition, the lack of an endoskeleton and a spinal cord precludes their use as models to study ageing in the bone systems or in the spinal cord (Ethan and McGinnis, 2004; Tissenbaum, 2015). For these reasons, vertebrate model organisms, which are more physiologically similar to humans, are invaluable model organisms to study a wider range of ageing-specific phenotypes that affect humans.

Vertebrate models

Zebrafish (*Danio rerio*) has become a tremendously successful vertebrate experimental model organism, and it is currently widely

used by a large scientific community. Zebrafish have transparent embryos, making them particularly amenable to live imaging studies in the early stages of development. Zebrafish have lower maintenance costs compared to mice, produce many embryos, and are amenable to large-scale genetic and pharmacological interventions. Based on these advantages, research on zebrafish has provided a fundamental contribution to the basic understanding of the molecular mechanisms underlying vertebrate development (Duboc et al., 2015; Kikuchi, 2015; Veldman and Lin, 2008). In addition, owing to their capacity of regenerating the heart, tail and spinal cord (Asnani and Peterson, 2014; Becker et al., 1997; Poss et al., 2003, 2002), they are also used as models for regenerative medicine (Goessling and North, 2014). These attributes have naturally paved the way for zebrafish to also be introduced as a model for studying ageing (Anchelin et al., 2011; Gilbert et al., 2014; Kishi, 2011; Kishi et al., 2003; Van Houcke et al., 2015). The model displays key hallmarks of ageing, including age-dependent mitochondrial dysfunction, telomere deterioration (loss of the terminal portions of the chromosomes) and protein oxidation (Kishi et al., 2003). Telomerase-deficient transgenic zebrafish (Anchelin et al., 2013; Henriques et al., 2013) (with a median lifespan of 9 months) show accelerated ageing phenotypes, demonstrating the important role of telomere length and stability in regulating vertebrate ageing and lifespan.

The laboratory mouse, *Mus musculus*, is the most widely adopted model system to investigate the biology of mammalian ageing. Given that the basic physiological mechanisms are highly conserved between mice and humans, the laboratory mouse has helped reveal many of the causal molecular mechanisms connecting ageing and ageing-related diseases (Liao and Kennedy, 2014; Vanhooren and Libert, 2013). The mouse ageing phenome includes up to 32 available inbred strains (Yuan et al., 2011), and the lifespan of the available inbred mouse strains varies from 2 to 4 years. The most commonly used strain in ageing research, C57BL/6, has a median lifespan of 914 days (Yuan et al., 2011). Such a short lifespan for a mammal makes it possible to test, in a relatively short time, the effects of any genetic, pharmacological or environmental intervention on mammalian ageing and lifespan. However, maintenance costs for mice are much higher compared to other model animals used to study ageing (Lieschke and Currie, 2007), and its lower litter size can limit opportunities for high-throughput screening under variable genetic, environmental and pharmacological interventions.

Importantly, the ageing vertebrate experimental model organisms listed above are characterised by a much longer lifespan than the invertebrate model organisms, substantially impacting experimental duration and costs. Therefore, there is a need to develop new model systems that share the advantages of short-lived non-vertebrate model organisms but demonstrate the physiological proximity to humans of vertebrate model organisms.

The African turquoise killifish (*Nothobranchius furzeri*), a teleost fish with a natural lifespan ranging between 4 and 9 months, is emerging as a new promising model organism in ageing research (Genade et al., 2005; Terzibasi et al., 2008). Below, we summarise the major advantages that make this species a competitive new model organism for ageing research. We describe its development, life cycle, known ageing phenotypes and the available approaches for modulating its lifespan experimentally, as well as the recent advances in transgenesis techniques and their applications in this animal. Finally, we discuss possible future applications of the turquoise killifish to understanding human ageing and ageing-associated diseases.

Turquoise killifish in nature and in the laboratory

In nature, the turquoise killifish dwells in seasonal water bodies in Mozambique and Zimbabwe (Reichard et al., 2009). Its habitat is yearly characterised by a brief rainy season followed by a longer dry season. During the rainy season, ephemeral ponds form along seasonal river drainages. The fish then rapidly hatch, reach sexual maturity in less than a month and reproduce before the water completely dries out in the subsequent dry season. The embryos are uniquely adapted to survive and develop in dry mud during the dry season (Blazek et al., 2013; Furness et al., 2015).

The current turquoise killifish laboratory strains include the inbred 'GRZ' strain, derived from an original population collected in 1968 (Genade et al., 2005; Valdesalici and Cellerino, 2003), and several wild strains that were derived more recently (Bartakova et al., 2013; Reichwald et al., 2009; Terzibasi et al., 2008). The GRZ strain has the shortest recorded lifespan among all the available turquoise killifish strains, with a median lifespan ranging from 9 to 16 weeks depending on the culture conditions (Genade et al., 2005; Kirschner et al., 2012; Terzibasi et al., 2008) (Fig. 1A, left), whereas longer-lived strains have median lifespans ranging from 23 to 28 weeks (Kirschner et al., 2012; Terzibasi et al., 2008; Valenzano et al., 2015) (Fig. 1A, right). Interestingly, although, in captivity, no difference in killifish survival is observed between the sexes (Valenzano et al., 2015, 2006b), large differences in sex ratios are observed in the wild, where females tend to be more frequent than males, which is also observed to some extent in other species, including humans (Reichard et al., 2009).

Life cycle of the turquoise killifish

The life cycle of the turquoise killifish is relatively unique because they are adapted to reach sexual maturity and reproduce during a very short (wet) period (Cellerino et al., 2015). Once hatched, fish grow very rapidly, as they need to complete sexual maturation and reproduce before the water completely evaporates (Podrabsky, 1999) (Fig. 1B). Each captive turquoise killifish female lays 20-40 eggs per day, with a maximum recorded number of 200 (Blazek et al., 2013; Graf et al., 2010; Polacik and Reichard, 2011), which is a similar order of magnitude as zebrafish (Eaton and Farley, 1974a,b; Hisaoka and Firlit, 1962; Kalueff et al., 2014), although zebrafish reproduce for a longer period. Although zebrafish and killifish can be bred in similar ways, and their embryo size is comparable, embryonic development in the turquoise killifish is slower (~2-3 weeks) than in zebrafish (~2-3 days) (Dolfi et al., 2014; Kimmel et al., 1995). However, owing to their rapid sexual maturation, which in killifish takes 3-4 weeks in captivity (Fig. 1B), the complete life cycle under controlled laboratory conditions is faster in the turquoise killifish (5.5-8 weeks) than in zebrafish (12-13 weeks) (<https://zfin.org/>) (Avdesh et al., 2012). The embryo development in the turquoise killifish is characterised by a developmentally arrested state called diapause (Wourms, 1972), in which embryos can survive over several months in the absence of water, encased in dry mud (Reichard et al., 2009). This special developmental state is functionally analogous to the *C. elegans* dauer state. Under controlled laboratory conditions, killifish embryos can skip diapause and therefore develop rapidly (Furness et al., 2015; Valenzano et al., 2011). Maternal age and temperature of embryo incubation are important factors regulating the exit from diapause in the killifish (Markofsky and Matias, 1977; Podrabsky et al., 2010). However, the molecular mechanisms involved in entry, persistence and exit from diapause are still largely uncharacterised in killifish. Because the genes controlling larval dauer in *C. elegans* have been shown to be




Fig. 1. Captive strains and the life cycle of the turquoise killifish. (A) Commonly used laboratory strains of the turquoise killifish: the short-lived strain (left; GRZ, yellow-tailed male fish) and long-lived strain (right; MZM-0403, red-tailed male fish). (B) Turquoise killifish life cycle (the time scale is based on the short-lived laboratory strain). Embryos can enter normal development or a developmentally arrested state called diapause, which lasts from a few weeks to several months and protects killifish during the dry season in the wild. Diapause consists of three different stages called diapause I, II and III. During the wet season in the wild (see main text) – and in laboratory conditions – hatched fry fully develop within 3-4 weeks and start spawning. Male fish are larger than females and have colourful fins and body, whereas the female fish are dull. Upon ageing ('old'), fish lose body colour, fin structure deteriorates and the spine becomes bent. The age for young, young adult and old fish is indicated in weeks.

crucial in regulating adult phenotypes, including ageing, it is important to identify what genes regulate killifish diapause and test whether they can also play a role in regulating adult physiology and ageing phenotypes, including ageing-related diseases. If the genes regulating killifish diapause also regulate adult phenotypes, including ageing, they could be relevant for understanding ageing-related phenotypes in humans.

Ageing phenotypes in the turquoise killifish

Despite its relatively short lifespan for a vertebrate, the turquoise killifish shows many molecular, cellular and physiological ageing phenotypes that are shared with many other organisms, including humans (Genade et al., 2005; Hartmann et al., 2009, 2011; Terzibasi et al., 2009, 2007; Valenzano et al., 2006b). Similarly to ageing mammals, who progressively lose hair and skin pigment with age (Geyfman and Andersen, 2010), male turquoise killifish – which are more colourful than females – progressively lose body and tail colour as well as their distinct patterning as they age (Fig. 1B). Old age in this short-lived vertebrate is also associated with abnormal spine curvature, defective vision, fin structure deterioration, decreased spontaneous locomotion activity, learning impairment (Genade et al., 2005; Valenzano et al., 2006b) and, interestingly, an increased risk of cancer (Baumgart et al., 2015). Fecundity also declines with age in the turquoise killifish, with embryo production reaching a peak at around 8-10 weeks and gradually declining thereafter (Blazek et al., 2013). These

macroscopic phenotypes recapitulate several of the complex age-dependent changes that occur in other vertebrates, including mouse and humans (Vanhooren and Libert, 2013). Compared to other fish, the striking feature of killifish ageing is its rapid onset within 3-4 months of age. For comparison, in zebrafish studies, individuals over 24 months of age are considered by researchers working on zebrafish models of ageing to be 'old' (Kishi et al., 2003).

Several ageing biomarkers have been developed to characterise the physiological age of killifish. Lipofuscin, a yellow-brown autofluorescent pigment whose concentration increases with age in several species, including humans (Goyal, 1982), accumulates in the brain and liver of old killifish (Goyal, 1982; Terzibasi et al., 2008, 2007). Senescence-associated β -galactosidase (SA- β -gal) staining, a marker for cellular senescence and stress response in human cells (Cristofalo, 2005; Dimri et al., 1995; Kurz et al., 2000; Untergasser et al., 2003; Yegorov et al., 1998), significantly increases in the skin of aged fish (Terzibasi et al., 2007). Neurodegeneration – measured by Fluoro-Jade B, which stains cell bodies, dendrites and axons of degenerating neurons but not those of healthy neurons (Schmued and Hopkins, 2000) – increases in fish brains from as early as 2 months of age, strongly suggesting a spontaneous age-dependent increase in neurodegeneration (Terzibasi et al., 2007; Valenzano et al., 2006b). The availability of various ageing biomarkers for the turquoise killifish allows characterisation of age-related changes in many tissues and under

different experimental conditions, further facilitating the use of this model system for ageing research.

Neoplastic lesions have been measured in turquoise killifish strains using several tumour-associated proteins, including Bcl-2, cytokeratin-8, carcinoembryonic antigen and mutated p53 (Di Cicco et al., 2011). In this study, the liver showed the highest rate of tumour incidence, and the short-lived GRZ strain showed an earlier onset of liver tumours compared to longer-lived strains. Additionally, the occurrence of these lesions increases as the fish aged (Di Cicco et al., 2011; Pompei et al., 2001), which is also observed in humans (de Magalhaes, 2013). Finally, liver cancer in killifish has a higher incidence in males than in females (Di Cicco et al., 2011), and this also replicates the pattern of liver cancer incidence in humans (Nordenstedt et al., 2010).

A range of factors could underlie liver cancer development in killifish, including chronic liver infection, metabolic perturbations (e.g. in carbohydrate metabolism) or genetic predisposition. It is of pivotal importance to compare cancer incidence in captive-bred and wild killifish populations to evaluate the relative contributions of environmental and genetic factors to this phenotype. Although liver cancer is not among the most common cancers in humans, it is one of the leading causes of cancer-related deaths worldwide (globocan.iarc.fr/Pages/fact_sheets_population.aspx). There are several mouse models of hepatocellular carcinoma, in which the condition can be induced chemically or genetically (Bakiri and Wagner, 2013). The distribution of the lesions and neoplasias in aged killifish, as gleaned from post-mortem analyses, largely differs from that in humans, mice and zebrafish (Di Cicco et al., 2011), indicating that mortality in killifish might be driven by cancer more than it is in other, longer-lived organisms. This, together with the fact that there are relatively few good models for these types of cancer in humans, highlight the potential to develop the turquoise killifish as a complementary model for liver cancer.

Age-dependent telomere attrition has been linked to organismal ageing in humans and in many other model organisms (Benetos et al., 2001; Harley et al., 1990; Lopez-Otin et al., 2013). Interestingly, killifish telomeres, which are over four times shorter than in mice (Zhu et al., 1998), are comparable in length to human telomeres (Hartmann et al., 2009). Although telomeres have been shown to shorten with age in a long-lived turquoise killifish strain, they did not shorten significantly in the short-lived GRZ strain, possibly owing to the very short lifespan of this strain (Hartmann et al., 2009). This suggests that telomere attrition might not contribute to rapid ageing of the short-lived GRZ strain. However, a telomerase mutant line of the turquoise killifish showed premature infertility, a dramatic decrease of red and white blood cell numbers, abnormalities in the epithelial cells of the intestine, including decreased polarity, and increased nuclear/cytoplasmic ratio (Harel et al., 2015). These results suggest that telomerase plays a key role in the maintenance of organismal homeostasis in the turquoise killifish, as in other organisms.

Mitochondrial DNA (mtDNA) instability has been associated with ageing in many species, including humans (Barazzoni et al., 2000; Lopez-Otin et al., 2013; Tauchi and Sato, 1968; Yen et al., 1989; Yui et al., 2003). mtDNA lacks protective histones; therefore, it is more likely to accumulate mutations compared to nuclear DNA (Tatarenkov and Avise, 2007). mtDNA mutations and deletions increase with age, whereas mtDNA copy number decreases, especially in the liver of mouse, rat and humans (Barazzoni et al., 2000; Bratic and Larsson, 2013). The increased mtDNA mutations and deletions with age drive an early onset of age-associated phenotypes (Vermulst et al., 2008). In line with this, mtDNA copy

number was found to be significantly reduced in several turquoise killifish tissues, including brain, liver and muscle, and its biogenesis was also impaired in muscles from old individuals (Hartmann et al., 2011).

The capacity to regenerate and repair tissues and organs helps to ensure homeostasis, resulting in the maintenance of an optimal health status. Interestingly, a recent study has shown that the capacity to regenerate the caudal fin after amputation was progressively impaired throughout ageing in the long-lived MZM-0703 turquoise killifish strain (Wendler et al., 2015). Whereas 8-week-old fish are capable of regenerating the caudal fin almost completely (98%) within 4 weeks, fish at 54 weeks of age could regenerate only 46% of the original fin size (Wendler et al., 2015).

Experimental modulation of the lifespan of turquoise killifish

Ageing is a highly integrated process that can be experimentally modulated by various types of interventions, including changes in nutrients, drugs, temperature and social conditions, as well as genetic modifications.

As mentioned above, DR can effectively modulate lifespan and ageing in several organisms, ranging broadly from yeast to worms, flies, beetles, chicken, mice, rats, dogs and macaques (Fontana and Partridge, 2015; Taormina and Mirisola, 2014). The effects of DR on lifespan and ageing biomarkers were tested in the turquoise killifish, which were fed every other day instead of daily (Terzibasi et al., 2009). The results were strain-dependent: DR resulted in prolonged lifespan in the short-lived GRZ strain but not in a wild-derived, long-lived MZM-0410 strain (Terzibasi et al., 2008). Under the DR regimen, the short-lived strain showed reduced neurodegeneration, slower accumulation of lipofuscin, improved learning performance and decreased occurrence of tumours (Terzibasi et al., 2009). This indicates that DR via every-other-day feeding can delay ageing in the turquoise killifish, depending on the genetic background. The effects of DR need testing under different nutrient restriction paradigms, including overall reduction of daily nutrient intake, or by varying the contribution of different micronutrients in the diet, which has shown to be effective in modulating ageing and longevity in other model organisms (Mair et al., 2005; Miller et al., 2005; Skorupa et al., 2008; Solon-Biet et al., 2015; Zimmerman et al., 2003). The effects of different nutrient restriction paradigms on humans are still largely unknown, warranting further investigation in the context of model systems, including the turquoise killifish.

Temperature is an important factor that has a huge effect in modulating metabolic rate and organism physiology (Conti, 2008; Lithgow, 1996). Modulating both environmental and individual temperature has a significant impact on organism physiology and can, within a specific range, modulate lifespan and ageing in many model organisms (Conti et al., 2006; Lamb, 1968; Liu and Walford, 1966; Mair et al., 2003). Decreased water temperature is sufficient to extend both the median (1 week) and maximum (1.5 weeks) lifespan of turquoise killifish, and leads to a 40% decrease in adult size compared to adult fish grown under regular culturing temperature, indicating a dramatic influence of temperature on metabolism. Several age-associated phenotypes, such as lipofuscin accumulation, spontaneous locomotor activity and learning performance, are also significantly improved in fish cultured at a lower temperature (Valenzano et al., 2006a). Importantly, previous studies performed in fish (Atlantic salmon and *Cynolebias adloffii*) showed that temperature changes can modulate lifespan and growth in different directions, and that there exist temperature optima for lifespan extension and growth optimisation (Handelanda et al.,

2008; Liu and Walford, 1966). However, studies that accurately correlate a broad range of temperatures with fish growth rates, survival and reproduction are still lacking, and these are needed to establish a clear mechanistic connection between environmental temperature, metabolism and life history. The understanding of the molecular mechanisms responsible for lifespan extension through temperature modulation could be used in the future to design molecular interventions that slow down ageing, retard the onset of age-related pathologies and ultimately extend lifespan.

The use of the natural polyphenol resveratrol, known to increase lifespan and delay ageing in worms and flies, can increase median and maximum lifespan in a dose-dependent manner in both male and female turquoise killifish. Compared to control-fed fish, resveratrol-fed fish remained physically active for a longer time, indicating that this compound is sufficient to retard the age-dependent decline in physical activity. Similarly, resveratrol-fed fish showed better learning performance at later ages than control-fed

fish (Genade and Lang, 2013; Liu et al., 2015; Valenzano and Cellierino, 2006; Valenzano et al., 2006b; Yu and Li, 2012). These effects are consistent with the biological effect of resveratrol on ageing and age-associated physiology in yeast, worms, flies and mice fed a high-fat diet (Baur et al., 2006; Marchal et al., 2013). Studies linking the effects of resveratrol on human metabolism and ageing are to date inconclusive and more work is needed to clarify the effects of this natural compound in our species.

The modulation of the turquoise killifish lifespan and ageing through external interventions such as diet, temperature and chemicals supports the use of this organism as an experimental platform for large-scale screens of age-modulating genes and chemicals.

Genetic modifications in the turquoise killifish

To date, two methods have been successfully developed to modify the turquoise killifish genome: random genome integration through the Tol2 DNA transposase and targeted genome editing using CRISPR/




Fig. 2. Side-by-side comparison of timing of transgenic line generation using genetic manipulations in the turquoise killifish, zebrafish and mouse. Synthesised single guide RNA (sgRNA) and Cas9 nucleases are injected into one-cell-stage embryos. Injected embryos are called F0 embryos. After hatching, transgenic fish are backcrossed to wild-type fish and generate F1 offspring. Further backcrosses are used to remove off-target mutations. Data are from the following references: turquoise killifish (Harel et al., 2015); zebrafish (Hwang et al., 2013; Jao et al., 2013); mouse (Wang et al., 2013).

Cas9 nuclease. To elicit genome editing, both methods require microinjection of RNA and DNA combinations into the one-cell-stage embryo (Table 1 and Fig. 2) (Harel et al., 2015; Hartmann and Englert, 2012; Valenzano et al., 2011). Whereas the Tol2 system was initially applied as a proof-of-principle to validate the possibility of generating transgenic turquoise killifish, CRISPR/Cas9 has recently been established in this species to directly test the role of genes involved in key ageing regulatory pathways (Harel et al., 2015). Deletion mutants or variants in relevant ageing pathways, including telomere attrition, deregulation of nutrient sensing, loss of proteostasis, genomic instability, mitochondrial dysfunction, epigenetic alteration, altered intercellular communication, cellular senescence and stem cell exhaustion, have been generated (Harel et al., 2015). At each cell division, telomeres cannot be fully replicated, and consequently shorten. This phenomenon of telomere deterioration progresses with ageing. Importantly, the loss of telomere length maintenance is crucial for cancer and degenerative diseases (Aubert and Lansdorp, 2008; Lopez-Otin et al., 2013). Telomerase reverse transcriptase (TERT) is a crucial protein whose key role is to elongate telomeres by adding nucleic acids (Aubert and Lansdorp, 2008). In humans, telomere dysfunction induces many degenerative diseases, including dyskeratosis congenita and pulmonary fibrosis. Harel et al. generated several *TERT* mutants in the turquoise killifish using the CRISPR/Cas9 nuclease system, and the deletion mutants lacking the catalytic function of TERT underwent age-dependent telomere shortening (Harel et al., 2015). In addition, *TERT* mutant fish

developed atrophied gonads and severe age-related morphological defects in actively proliferating tissues such as testes, intestine and blood, compared to less actively proliferating tissues such as heart, muscle, liver and kidney (Harel et al., 2015). The non-synonymous variant of *TERT* corresponding to human dyskeratosis congenita was successfully generated by homology-directed repair accompanied by the CRISPR/Cas9 nuclease system, showing that this strategy can be effectively used to generate fish models of human-specific diseases (Harel et al., 2015).

As well as engineering the genome of turquoise killifish, direct injection of synthetic RNA into the one-cell-stage embryo recently enabled the development of a turquoise killifish fluorescence ubiquitination cell cycle indicator (FUCCI) (Dolfi et al., 2014). This method takes advantage of two fluorescent-tagged proteins that differentially degrade during the cell cycle, resulting in the accumulation of red fluorescence at the G1 phase of the cell cycle and of green fluorescence at the S/G2/M phases (Sakaue-Sawano et al., 2008). FUCCI expression remains stable for 3-4 days after the FUCCI mRNA injection into killifish embryos and allows the temporal tracking of early cell-division kinetics. When stable FUCCI transgenic killifish lines become available, this tool will become important in monitoring cellular proliferation during development, tumour formation, tissue/organ regeneration and senescence.

Genetic modifications in the turquoise killifish are rapid and highly efficient, permitting generation of stable transgenic lines more rapidly than in any other available vertebrate model (Fig. 2).

Table 2. Turquoise killifish as a platform to test gene variants associated with human age-related dysfunctions

Dysfunctions in human	Disease examples	Key genes in human	Killifish orthologues
Genome instability	Cancer	<i>P53, PTEN, PI3K, HER2, VEGFR, PARP</i>	<i>PTEN, PARP</i>
	Aplastic anaemia	<i>TERT, TERC, TRF1/2</i>	<i>TERT</i>
	Dyskeratosis congenita	<i>TERT, TERC, DKC</i>	<i>TERT, DKC1</i>
	Werner syndrome	<i>WRN</i>	<i>WRN</i>
	Progeria	<i>LMNA</i>	<i>LMNA</i>
Mitochondrial dysfunction	Alpers-Huttenlocher syndrome	<i>POLG</i>	<i>POLG</i>
	Ataxia neuropathy syndromes	<i>POLG, C10orf2</i>	<i>POLG</i>
	Leigh syndrome	<i>MT-ATP6, SURF1</i>	<i>ATP6</i>
	Neuropathy, ataxia, and retinitis pigmentosa	<i>MT-ATP6</i>	<i>ATP6</i>
Neurodegeneration	Alzheimer's disease	<i>APP, PSEN1/2, APOE, TREM2</i>	<i>APP, PSEN1/2</i>
	Huntington's disease	<i>HTT</i>	<i>HTT</i>
	Parkinson's disease	<i>LRRK2, PINK1, SNCA, FBXO7, PARK2, PARK7/DJ-1, PLA2G6, VPS35, ATP13A2, DNAJC6, SYNJ1</i>	<i>LRRK2, PINK1, SCNA, FOXO7, PARK2, VPS35, PARK7, PLA2G6, ATP13A2, DNAJC6, SYNJ1</i>
Metabolic dysfunction	Phenylketonuria	<i>PAH</i>	<i>PAH</i>
	Propionic acidemia	<i>PCCA, PCCB</i>	<i>PCCA, PCCB</i>
	Glycogen storage diseases	<i>G6PC, SLC37A4, GYS1/2, PYGL, PYGM</i>	<i>G6PC, SLC37A4, GYS1/2, PYGL, PYGM</i>
	Tay-Sachs disease Type 1 diabetes	<i>HEXA</i> <i>HLA-DQA1, HLA-DQB1, HLA-DRB1</i>	<i>HEXA</i> –
Autoimmune defects	Systemic lupus erythematosus	<i>HLA-A/B/C, HLA-DP/DQ/DR</i>	<i>HLA-DPA1, HLA-DPB1, HLA-DQB2</i>
	Rheumatoid arthritis	<i>HLA-DRB1</i>	–
	Multiple sclerosis	<i>HLA-DRB1, IL-7R</i>	<i>IL7R</i>
Cardiovascular dysfunction	Wolff-Parkinson-White syndrome	<i>PRKAG2</i>	<i>PRKAG2A/B</i>
	Progressive familial heart block	<i>SCN5A, TRPM4</i>	<i>SCN5A</i>
	McKusick-Kaufman syndrome	<i>MKKS</i>	<i>MKKS</i>

Combined with their other advantageous features, as described above, efficient genetic manipulation of the turquoise killifish could allow it to become a powerful and highly scalable platform to study age-related changes and diseases (Table 2).

Looking forward: potential applications and limitations of killifish models of ageing

The long lifespan of the currently available vertebrate model organisms is a major limitation for the feasibility of large-scale ageing studies. The development of new model organisms is vital to compensate for the limitations of previous model systems. In the past, several fish species have been adopted as potential ageing model organisms, including medaka (Ding et al., 2010; Egami and Eto, 1973; Gopalakrishnan et al., 2013), guppies (Reznick et al., 2006; Woodhead and Pond, 1984; Woodhead et al., 1983) and both American as well as African killifish (Liu and Walford, 1966; Markofsky and Milstoc, 1979a,b). As highlighted in this article, the turquoise killifish has emerged as a promising new model organism for vertebrate ageing because it uniquely combines a short lifespan and life cycle with vertebrate-specific features that are missing from the currently used non-vertebrate model organisms. In particular, it undergoes continuous adult cellular proliferation, has adult stem cells in many tissues and an adaptive immune system, and shows a spontaneous age-dependent increased risk for cancer. The turquoise killifish has the shortest lifespan among all vertebrate models in captivity, and shares several age-associated phenotypes with other vertebrates, including humans (Table 1). Recent genomic and transcriptome data analysis in the turquoise killifish have revealed many orthologous genes to humans and other model organisms (Harel et al., 2015; Petzold et al., 2013; Reichwald et al., 2009). Additionally, the fully sequenced, assembled and annotated genome is now available, together with a high-density Rad-Seq genome map, as well as ChipSeq and transcriptome data from several tissues (Reichwald et al., 2015; Valenzano et al., 2015). The availability of these genomic tools will enable a broad scientific community to take advantage of this model system in an unprecedented way. The turquoise killifish captive strains are highly fecund, facilitating transgenic line generation and genetic screenings (Fig. 2). Importantly, there exists a highly inbred turquoise killifish strain (GRZ) that enables easy testing of the effects of biochemical or genetic modifications on organismal physiology. Thus, the turquoise killifish is likely to provide a very useful platform for testing the function of genes involved in longevity regulation, particularly to examine the underlying genes of extreme longevity. Unique gene variants have been found in human centenarians, as well as very long-lived model organisms such as the naked mole rat and bowhead whale (Keane et al., 2015; Kim et al., 2011; Sebastiani et al., 2011; Willcox et al., 2006). However, their causal role in ageing and longevity is largely unknown. Given the long lifespan of mice and zebrafish, it is impractical to functionally analyse such gene variants, especially when the expected outcome is lifespan extension. The turquoise killifish could fill the need for a short-lived vertebrate, enabling variants linked to extreme longevity to be tested in a rapid and effective way. In parallel, the turquoise killifish allows rapid testing – in vertebrates – of the genetic variants identified in worms and flies (de Magalhaes and Costa, 2009; Tacutu et al., 2013), enabling the translation of findings from short-lived invertebrates to vertebrates.

Human ageing is associated with an increased risk of several diseases, such as cancer or Alzheimer's disease (Table 2), which have a strong genetic component. The turquoise killifish provides the opportunity to test, in a short time, the causal role of specific gene variants in the onset of age-associated diseases. For instance, to

assess the effects of a given genetic manipulation on lifespan would be six times faster in killifish than in mice (Fig. 2).

Outbred turquoise killifish strains are used to identify the genetic architecture of naturally occurring phenotypes that differ among strains, such as male colouration, susceptibility to pigment aberration, and survival. Quantitative trait locus (QTL) mapping studies in the turquoise killifish have revealed genomic loci that are significantly associated with such phenotypes (Kirschner et al., 2012; Valenzano et al., 2009). Previous studies in sticklebacks have used genetic mapping to identify the genetic basis of human phenotypic variation (Miller et al., 2007). The possibility to combine genetic mapping with transgenesis in the turquoise killifish could help reveal the basic molecular mechanisms underlying the susceptibility to early ageing and age-related diseases, which might be shared with humans.

Reverse genetic tools, which use transposases and nuclease-based systems to introduce genomic integrations and deletions, were recently developed in the turquoise killifish (Allard et al., 2013; Harel et al., 2015; Hartmann and Englert, 2012; Valenzano et al., 2011). However, forward genetic approaches have not been explicitly tested in the turquoise killifish yet. However, the use of the Tol2 transposase, which introduces portions of DNA in random regions of the host genome – similar to insertional mutagenesis – demonstrates the feasibility of forward genetic applications in this species (Valenzano et al., 2011). Random mutagenesis, using various mutagens such as N-ethyl-N-nitrosourea (ENU), has been used for a long time in many species – from bacteria to yeast, plants and animals – as a powerful method to stably alter the genetic information of an organism, and then to subsequently identify a causal connection between genetic modification and a specific biological phenotype (Muller, 1927), including longevity (Caspary, 2010; Hardy et al., 2010; Jorgensen and Mango, 2002). Several strategies for large-scale screening after chemical-induced mutagenesis in zebrafish have been developed (for a review, see Patton and Zon, 2001), which have resulted in a vast collection of zebrafish mutant lines, each having a specifically altered phenotype (Driever et al., 1996; Haffter et al., 1996; Lawson and Wolfe, 2011; Wang et al., 2012; Westerfield, 2000). Owing to the relevance of the turquoise killifish to study adult phenotypes, mutagenised fish should be screened for the phenotype of interest after sexual maturation. Unlike embryo or larval screens, which can take advantage of high population densities of individuals and rapid experimental time, adult screens – in particular those aimed at selecting longer-lived individuals – require a larger space and take longer. Even though turquoise killifish males can be territorial – especially when grown in isolation – this species allows for survival screens in group-housed conditions. Experiments to establish the effect of fish population density on experimental survival are currently ongoing in the Valenzano lab. However, the application of random mutagenesis to screen for turquoise killifish mutants that live longer requires large breeding facilities. In fact, screening for longevity mutants requires that all of the mutagenised fish have to breed, since by the time they breed it is not possible to assess whether they will be long-lived or short-lived. Only the offspring of the long-lived mutants will be used to generate mutant fish lines. An alternative strategy, widely employed in the *C. elegans* and *D. melanogaster* fields to identify genes involved in the regulation of longevity, is to induce mutations and then screen juvenile fish for stress resistance against chemicals or physical stressors (Denzel et al., 2014; Lin et al., 1998; Walker et al., 1998). This alternative strategy might help to overcome spatial and temporal limitations of longevity screening using exclusively adult turquoise killifish. The application of random mutagenesis accompanied by accessible

phenotypic analyses of the turquoise killifish has not yet been developed but has great potential to reveal currently unknown molecular mechanisms of ageing and age-associated diseases. The turquoise killifish also has the potential to be used for high-throughput drug screening in a similar way to zebrafish (Gibert et al., 2013), with the advantage of a shorter lifespan and life cycle. This application would make it an ideal system to test the effects of drugs on adult-specific phenotypes involved in ageing, age-related diseases and overall longevity.

Although model organisms in ageing research have greatly improved our understanding of the shared features of the ageing process across organisms, no single model is a perfect model of human ageing. It is important to note the intrinsic limitations of the turquoise killifish. For example, it cannot be directly used to study organs that are not shared between fish and mammals, such as lungs or bone marrow. Additionally, being a novel model organism, it still lacks important community-based research resources, such as a large-scale stock centre with wild-derived, transgenic and mutant strains, which are readily available in other model organisms that are more established. However, molecular tools are rapidly developing in the turquoise killifish, including several linkage maps (Kirschner et al., 2012; Valenzano et al., 2015, 2009), a transcriptome atlas (Petzold et al., 2013), and a completely sequenced, assembled and annotated genome (Reichwald et al., 2015; Valenzano et al., 2015). The short lifespan of this species, one of its biggest strengths as a vertebrate model organism, could potentially also constitute a limitation for its applications as a 'natural ageing' model system, because some of the very late-age phenotypes occurring for instance in humans (e.g. cardiovascular disease and dementia) might be not shared in this species. There are, however, several strains within this species, some of which differ substantially in captive lifespan, and a comparative study of age-dependent phenotypes in multiple strains could be used as an approach to investigate a larger range of such phenotypes. Additionally, other species of the same genus, which are longer-lived than the turquoise killifish, are available and can be housed under conditions similar to the turquoise killifish (Genade et al., 2005; Terzibasi et al., 2008), offering an additional opportunity for the comparative study of the molecular and genetic basis of vertebrate ageing.

In summary, the short-lived turquoise killifish has the potential to become a central model system in the field of the biology of ageing and be used as a bridge from the short-lived invertebrates to the longer-lived mammalian models in the study of the biological mechanisms that underlie age-associated diseases (Table 2). The growing scientific community working on the turquoise killifish is constantly developing novel genetic tools and a wide set of resources, which we predict will build this model up as a strong platform for the discovery of novel basic mechanisms that play an important role in vertebrate ageing.

Acknowledgements

We thank Dr S. J. Lee, the members of the Valenzano lab at the Max Planck Institute for Biology of Ageing, two anonymous reviewers and DMM's Reviews Editor, Paraminder Dhillon, for their constructive feedback on the manuscript.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Y.K. and D.R.V. conceived and wrote the Review, and H.G.N. significantly contributed to the text.

Funding

This work has been supported by the Max Planck Society, by the Max Planck Institute for Biology of Ageing in Cologne, Germany, and by the Research Center Program of Institute for Basic Science in the Republic of Korea (IBS-R013-D1).

References

- Akushевич, I., Kravchenko, J., Ukrantseva, S., Arbeev, K. and Yashin, A. I. (2013). Time trends of incidence of age-associated diseases in the US elderly population: medicare-based analysis. *Age Ageing* **42**, 494-500.
- Allard, J. B., Kamei, H. and Duan, C. (2013). Inducible transgenic expression in the short-lived fish *Nothobranchius furzeri*. *J. Fish Biol.* **82**, 1733-1738.
- Almada-Pagan, P. F., Lucas-Sanchez, A. and Tocher, D. R. (2014). Changes in mitochondrial membrane composition and oxidative status during rapid growth, maturation and aging in zebrafish, *Danio rerio*. *Biochim. Biophys. Acta* **1841**, 1003-1011.
- Anchelin, M., Murcia, L., Alcaraz-Pérez, F., García-Navarro, E. M. and Cayuela, M. L. (2011). Behaviour of telomere and telomerase during aging and regeneration in zebrafish. *PLoS ONE* **6**, e16955.
- Anchelin, M., Alcaraz-Perez, F., Martinez, C. M., Bernabe-Garcia, M., Mulero, V. and Cayuela, M. L. (2013). Premature aging in telomerase-deficient zebrafish. *Dis. Model. Mech.* **6**, 1101-1112.
- Armstrong, D., Rinehart, R., Dixon, L. and Reigh, D. (1978). Changes of peroxidase with age in *Drosophila*. *Age* **1**, 8-12.
- Asnani, A. and Peterson, R. T. (2014). The zebrafish as a tool to identify novel therapies for human cardiovascular disease. *Dis. Model. Mech.* **7**, 763-767.
- Aubert, G. and Lansdorp, P. M. (2008). Telomeres and aging. *Physiol. Rev.* **88**, 557-579.
- Avdesh, A., Chen, M., Martin-Iverson, M. T., Mondal, A., Ong, D., Rainey-Smith, S., Taddei, K., Lardelli, M., Groth, D. M., Verdile, G. et al. (2012). Regular care and maintenance of a zebrafish (*Danio rerio*) laboratory: an introduction. *J. Vis. Exp.* **69**, e4196.
- Bailey, K. J., Maslov, A. Y. and Pruitt, S. C. (2004). Accumulation of mutations and somatic selection in aging neural stem/progenitor cells. *Aging Cell* **3**, 391-397.
- Bakiri, L. and Wagner, E. F. (2013). Mouse models for liver cancer. *Mol. Oncol.* **7**, 206-223.
- Barazzoni, R., Short, K. R. and Nair, K. S. (2000). Effects of aging on mitochondrial DNA copy number and cytochrome c oxidase gene expression in rat skeletal muscle, liver, and heart. *J. Biol. Chem.* **275**, 3343-3347.
- Bartakova, V., Reichard, M., Janko, K., Polacik, M., Blazek, R., Reichwald, K., Cellerino, A. and Bryja, J. (2013). Strong population genetic structuring in an annual fish, *Nothobranchius furzeri*, suggests multiple savannah refugia in southern Mozambique. *BMC Evol. Biol.* **13**, 196.
- Baumgart, M., Di Cicco, E., Rossi, G., Cellerino, A. and Tozzini, E. T. (2015). Comparison of captive lifespan, age-associated liver neoplasias and age-dependent gene expression between two annual fish species: *nothobranchius furzeri* and *Nothobranchius korthause*. *Biogerontology* **16**, 63-69.
- Baur, J. A., Pearson, K. J., Price, N. L., Jamieson, H. A., Lerin, C., Kalra, A., Prabhu, V. V., Allard, J. S., Lopez-Lluch, G., Lewis, K. et al. (2006). Resveratrol improves health and survival of mice on a high-calorie diet. *Nature* **444**, 337-342.
- Becker, T., Wullimann, M. F., Becker, C. G., Bernhardt, R. R. and Schachner, M. (1997). Axonal regrowth after spinal cord transection in adult zebrafish. *J. Comp. Neurol.* **377**, 577-595.
- Benetos, A., Okuda, K., Lajemi, M., Kimura, M., Thomas, F., Skurnick, J., Labat, C., Bean, K. and Aviv, A. (2001). Telomere length as an indicator of biological aging: the gender effect and relation with pulse pressure and pulse wave velocity. *Hypertension* **37**, 381-385.
- Blazek, R., Polacik, M. and Reichard, M. (2013). Rapid growth, early maturation and short generation time in African annual fishes. *Evodevo* **4**, 24.
- Bonawitz, N. D., Chatenay-Lapointe, M., Pan, Y. and Shadel, G. S. (2007). Reduced TOR signaling extends chronological life span via increased respiration and upregulation of mitochondrial gene expression. *Cell Metab.* **5**, 265-277.
- Bratic, A. and Larsson, N.-G. (2013). The role of mitochondria in aging. *J. Clin. Invest.* **123**, 951-957.
- Brenner, S. (1974). The genetics of *Caenorhabditis elegans*. *Genetics* **77**, 71-94.
- Brody, J. A. and Grant, M. D. (2001). Age-associated diseases and conditions: implications for decreasing late life morbidity. *Aging* **13**, 64-67.
- Burhans, W. C. and Weinberger, M. (2012). DNA damage and DNA replication stress in yeast models of aging. *Subcell Biochem.* **57**, 187-206.
- Burtner, C. R. and Kennedy, B. K. (2010). Progeria syndromes and ageing: what is the connection? *Nat. Rev. Mol. Cell Biol.* **11**, 567-578.
- Cagney, G. (2009). Interaction networks: lessons from large-scale studies in yeast. *Proteomics* **9**, 4799-4811.
- Caspary, T. (2010). Phenotype-driven mouse ENU mutagenesis screens. *Methods Enzymol.* **477**, 313-327.
- Cellerino, A., Valenzano, D. R. and Reichard, M. (2015). From the bush to the bench: the annual *Nothobranchius* fishes as a new model system in biology. *Biol. Rev. Camb. Philos. Soc.* doi:10.1111/brv.12183.
- Charlesworth, B. (2000). Fisher, Medawar, Hamilton and the evolution of aging. *Genetics* **156**, 927-931.
- Chintapalli, V. R., Wang, J. and Dow, J. A. T. (2007). Using FlyAtlas to identify better *Drosophila melanogaster* models of human disease. *Nat. Genet.* **39**, 715-720.
- Collins, J. J., Huang, C., Hughes, S. and Kornfeld, K. (2008). The measurement and analysis of age-related changes in *Caenorhabditis elegans*. *WormBook*, 1-21.

- Colman, R. J., Beasley, T. M., Kemnitz, J. W., Johnson, S. C., Weindruch, R. and Anderson, R. M. (2014). Caloric restriction reduces age-related and all-cause mortality in rhesus monkeys. *Nat. Commun.* **5**, 3557.
- Conti, B. (2008). Considerations on temperature, longevity and aging. *Cell Mol. Life Sci.* **65**, 1626-1630.
- Conti, B., Sanchez-Alavez, M., Winsky-Sommerer, R., Morale, M. C., Lucero, J., Brownell, S., Fabre, V., Huitron-Resendiz, S., Henriksen, S., Zorrilla, E. P. et al. (2006). Transgenic mice with a reduced core body temperature have an increased life span. *Science* **314**, 825-828.
- Cooper, A. A., Gitler, A. D., Cashikar, A., Haynes, C. M., Hill, K. J., Bhullar, B., Liu, K. N., Xu, K. X., Strathearn, K. E., Liu, F. et al. (2006). alpha-synuclein blocks ER-Golgi traffic and Rab1 rescues neuron loss in Parkinson's models. *Science* **313**, 324-328.
- Cota, D., Matter, E. K., Woods, S. C. and Seeley, R. J. (2008). The role of hypothalamic mammalian target of rapamycin complex 1 signaling in diet-induced obesity. *J. Neurosci.* **28**, 7202-7208.
- Craig, T., Smelick, C., Tacutu, R., Wuttke, D., Wood, S. H., Stanley, H., Janssens, G., Savitskaya, E., Moskalev, A., Arking, R. et al. (2015). The Digital Ageing Atlas: integrating the diversity of age-related changes into a unified resource. *Nucleic Acids Res.* **43**, D873-D878.
- Cristofalo, V. J. (2005). SA beta Gal staining: biomarker or delusion. *Exp. Gerontol.* **40**, 836-838.
- Croll, N. A., Smith, J. M. and Zuckerman, B. M. (1977). The aging process of the nematode *Caenorhabditis elegans* in bacterial and axenic culture. *Exp. Aging Res.* **3**, 175-189.
- Curtisinger, J. W., Fukui, H. H., Townsend, D. R. and Vaupel, J. W. (1992). Demography of genotypes: failure of the limited life-span paradigm in *Drosophila melanogaster*. *Science* **258**, 461-463.
- de Magalhaes, J. P. (2004). From cells to ageing: a review of models and mechanisms of cellular senescence and their impact on human ageing. *Exp. Cell Res.* **300**, 1-10.
- de Magalhaes, J. P. (2013). How ageing processes influence cancer. *Nat. Rev. Cancer* **13**, 357-365.
- de Magalhaes, J. P. and Costa, J. (2009). A database of vertebrate longevity records and their relation to other life-history traits. *J. Evol. Biol.* **22**, 1770-1774.
- De Sandre-Giovannoli, A., Bernard, R., Cau, P., Navarro, C., Amiel, J., Boccaccio, I., Lyonnet, S., Stewart, C. L., Munnich, A., Le Merrer, M. et al. (2003). Lamin A truncation in Hutchinson-Gilford progeria. *Science* **300**, 2055-2055.
- Demetrius, L. (2006). Aging in mouse and human systems: a comparative study. *Ann. N. Y. Acad. Sci.* **1067**, 66-82.
- Demontis, F. and Perrimon, N. (2010). FOXO/4E-BP signaling in *Drosophila* muscles regulates organism-wide proteostasis during aging. *Cell* **143**, 813-825.
- Denzel, M. S., Storm, N. J., Gutschmidt, A., Baddi, R., Hinze, Y., Jarosch, E., Sommer, T., Hoppe, T. and Antebi, A. (2014). Hexosamine pathway metabolites enhance protein quality control and prolong life. *Cell* **156**, 1167-1178.
- Di Cicco, E., Tozzini, E. T., Rossi, G. and Cellerino, A. (2011). The short-lived annual fish *Nothobranchius furzeri* shows a typical teleost aging process reinforced by high incidence of age-dependent neoplasias. *Exp. Gerontol.* **46**, 249-256.
- Dimri, G. P., Lee, X., Basile, G., Acosta, M., Scott, G., Roskelley, C., Medrano, E. E., Linskens, M., Rubelj, I., Pereira-Smith, O. et al. (1995). A biomarker that identifies senescent human cells in culture and in aging skin in vivo. *Proc. Natl. Acad. Sci. USA* **92**, 9363-9367.
- Ding, L., Kuhne, W. W., Hinton, D. E., Song, J. and Dynan, W. S. (2010). Quantifiable biomarkers of normal aging in the Japanese medaka fish (*Oryzias latipes*). *PLoS ONE* **5**, e13287.
- Dolfi, L., Ripa, R. and Cellerino, A. (2014). Transition to annual life history coincides with reduction in cell cycle speed during early cleavage in three independent clades of annual killifish. *Evodevo* **5**, 32.
- Dolle, M. E. T., Snyder, W. K., Gossen, J. A., Lohman, P. H. M. and Vijg, J. (2000). Distinct spectra of somatic mutations accumulated with age in mouse heart and small intestine. *Proc. Natl. Acad. Sci. USA* **97**, 8403-8408.
- Driever, W., Solnica-Krezel, L., Schier, A. F., Neuhauss, S. C., Malicki, J., Stemple, D. L., Stainier, D. Y., Zwartkruis, F., Abdelilah, S., Rangini, Z. et al. (1996). A genetic screen for mutations affecting embryogenesis in zebrafish. *Development* **123**, 37-46.
- Duboc, V., Dufourcq, P., Blader, P. and Roussigne, M. (2015). Asymmetry of the brain: development and implications. *Annu. Rev. Genet.* **49**, 647-672.
- Eaton, R. C. and Farley, R. D. (1974a). Growth and reduction of dispensation of zebrafish, *Brachydanio rerio*, in the laboratory. *Copeia* **1974**, 204-209.
- Eaton, R. C. and Farley, R. D. (1974b). Spawning cycle and egg production of zebrafish, *Brachydanio rerio*, in the Laboratory. *Copeia* **1974**, 195-204.
- Echigoya, Y., Aoki, Y., Miskew, B., Panesar, D., Touznik, A., Nagata, T., Tanihata, J., Nakamura, A., Nagaraju, K. and Yokota, T. (2015). Long-term efficacy of systemic multiexon skipping targeting dystrophin exons 45-55 with a cocktail of vivo-morpholinos in mdx52 mice. *Mol. Ther. Nucleic Acids* **4**, e225.
- Egami, N. and Eto, H. (1973). Effect of x-irradiation during embryonic stage in life span in the fish, *Oryzias latipes*. *Exp. Gerontol.* **8**, 219-222.
- Eisenstein, M. (2012). Centenarians: great expectations. *Nature* **492**, S6-S8.
- Ethan, B. and McGinnis, W. (2004). Model organisms in the study of development and disease. *Oxford Monogr. Med. Genet.* **49**, 25-45.
- Fahlstrom, A., Yu, Q. and Ulfhake, B. (2011). Behavioral changes in aging female C57BL/6 mice. *Neurobiol. Aging* **32**, 1868-1880.
- Ferguson, M., Mockett, R. J., Shen, Y., Orr, W. C. and Sohal, R. S. (2005). Age-associated decline in mitochondrial respiration and electron transport in *Drosophila melanogaster*. *Biochem. J.* **390**, 501-511.
- Fontana, L. and Partridge, L. (2015). Promoting health and longevity through diet: from model organisms to humans. *Cell* **161**, 106-118.
- Fraenkel, M., Ketzinel-Gilad, M., Ariav, Y., Pappo, O., Karaca, M., Castel, J., Berthault, M.-F., Magnan, C., Cerasi, E., Kaiser, N. et al. (2008). mTOR inhibition by rapamycin prevents beta-cell adaptation to hyperglycemia and exacerbates the metabolic state in type 2 diabetes. *Diabetes* **57**, 945-957.
- Furness, A. I., Lee, K. and Reznick, D. N. (2015). Adaptation in a variable environment: phenotypic plasticity and bet-hedging during egg diapause and hatching in an annual killifish. *Evolution* **69**, 1461-1475.
- Genade, T. and Lang, D. M. (2013). Resveratrol extends lifespan and preserves glia but not neurons of the *Nothobranchius guentheri* optic tectum. *Exp. Gerontol.* **48**, 202-212.
- Genade, T., Benedetti, M., Terzibas, E., Roncaglia, P., Valenzano, D. R., Cattaneo, A. and Cellerino, A. (2005). Annual fishes of the genus *Nothobranchius* as a model system for aging research. *Aging Cell* **4**, 223-233.
- Gerhard, G. S., Kauffman, E. J., Wang, X., Stewart, R., Moore, J. L., Kasales, C. J., Demidenko, E. and Cheng, K. C. (2002). Life spans and senescent phenotypes in two strains of Zebrafish (*Danio rerio*). *Exp. Gerontol.* **37**, 1055-1068.
- Geyfman, M. and Andersen, B. (2010). Clock genes, hair growth and aging. *Aging* **2**, 122-128.
- Gibert, Y., Trengove, M. C. and Ward, A. C. (2013). Zebrafish as a genetic model in pre-clinical drug testing and screening. *Curr. Med. Chem.* **20**, 2458-2466.
- Gilbert, M. J. H., Zerulla, T. C. and Tierney, K. B. (2014). Zebrafish (*Danio rerio*) as a model for the study of aging and exercise: physical ability and trainability decrease with age. *Exp. Gerontol.* **50**, 106-113.
- Girard, L. R., Fiedler, T. J., Harris, T. W., Carvalho, F., Antoshechkin, I., Han, M., Sternberg, P. W., Stein, L. D. and Chalfie, M. (2007). WormBook: the online review of *Caenorhabditis elegans* biology. *Nucleic Acids Res.* **35**, D472-D475.
- Goessling, W. and North, T. E. (2014). Repairing quite swimmingly: advances in regenerative medicine using zebrafish. *Dis. Model. Mech.* **7**, 769-776.
- Gopalakrishnan, S., Cheung, N. K. M., Yip, B. W. P. and Au, D. W. T. (2013). Medaka fish exhibits longevity gender gap, a natural drop in estrogen and telomere shortening during aging: a unique model for studying sex-dependent longevity. *Front. Zool.* **10**, 78.
- Gottlieb, S. and Ruvkun, G. (1994). daf-2, daf-16 and daf-23: genetically interacting genes controlling Dauer formation in *Caenorhabditis elegans*. *Genetics* **137**, 107-120.
- Goyal, V. K. (1982). Lipofuscin pigment accumulation in human brain during aging. *Exp. Gerontol.* **17**, 481-487.
- Graf, M., Cellerino, A. and Englert, C. (2010). Gender separation increases somatic growth in females but does not affect lifespan in *Nothobranchius furzeri*. *PLoS ONE* **5**, e11958.
- Grotewiel, M. S., Martin, I., Bhandari, P. and Cook-Wiens, E. (2005). Functional senescence in *Drosophila melanogaster*. *Ageing Res. Rev.* **4**, 372-397.
- Guo, L., Karpac, J., Tran, S. L. and Jasper, H. (2014). PGRP-SC2 promotes gut immune homeostasis to limit commensal dysbiosis and extend lifespan. *Cell* **156**, 109-122.
- Haddadi, M., Jahromi, S. R., Sagar, B. K. C., Patil, R. K., Shivanandappa, T. and Ramesh, S. R. (2014). Brain aging, memory impairment and oxidative stress: a study in *Drosophila melanogaster*. *Behav. Brain Res.* **259**, 60-69.
- Haffter, P., Granato, M., Brand, M., Mullins, M. C., Hammerschmidt, M., Kane, D. A., Odenthal, J., van Eeden, F. J., Jiang, Y. J., Heisenberg, C. P. et al. (1996). The identification of genes with unique and essential functions in the development of the zebrafish, *Danio rerio*. *Development* **123**, 1-36.
- Hahn, J. H., Kim, S., DiLoreto, R., Shi, C., Lee, S. J., Murphy, C. T. and Nam, H. G. (2015). *C. elegans* maximum velocity correlates with healthspan and is maintained in worms with an insulin receptor mutation. *Nat. Commun.* **6**, 8919.
- Hamilton, B., Dong, Y., Shindo, M., Liu, W., Odell, I., Ruvkun, G. and Lee, S. S. (2005). A systematic RNAi screen for longevity genes in *C. elegans*. *Genes Dev.* **19**, 1544-1555.
- Handelanda, S. O., Imsland, A. K. and Stefansson, S. O. (2008). The effect of temperature and fish size on growth, feed intake, food conversion efficiency and stomach evacuation rate of Atlantic salmon post-smolts. *Aquaculture* **283**, 36-42.
- Hardy, S., Legagneux, V., Audic, Y. and Paillard, L. (2010). Reverse genetics in eukaryotes. *Biol. Cell* **102**, 561-580.
- Harel, I., Benayoun, B. A., Machado, B., Singh, P. P., Hu, C.-K., Pech, M. F., Valenzano, D. R., Zhang, E., Sharp, S. C., Artandi, S. E. et al. (2015). A platform for rapid exploration of aging and diseases in a naturally short-lived vertebrate. *Cell* **160**, 1013-1026.
- Harley, C. B., Futcher, A. B. and Greider, C. W. (1990). Telomeres shorten during aging of human fibroblasts. *Nature* **345**, 458-460.

- Hartmann, N. and Englert, C. (2012). A microinjection protocol for the generation of transgenic killifish (Species: *Nothobranchius furzeri*). *Dev. Dyn.* **241**, 1133-1141.
- Hartmann, N., Reichwald, K., Lechel, A., Graf, M., Kirschner, J., Dorn, A., Terzibasi, E., Wellner, J., Platzer, M., Rudolph, K. L. et al. (2009). Telomeres shorten while Tert expression increases during ageing of the short-lived fish *Nothobranchius furzeri*. *Mech. Ageing Dev.* **130**, 290-296.
- Hartmann, N., Reichwald, K., Wittig, I., Drose, S., Schmeisser, S., Luck, C., Hahn, C., Graf, M., Gausmann, U., Terzibasi, E. et al. (2011). Mitochondrial DNA copy number and function decrease with age in the short-lived fish *Nothobranchius furzeri*. *Aging Cell* **10**, 824-831.
- Hashizume, O., Ohnishi, S., Mito, T., Shimizu, A., Iashikawa, K., Nakada, K., Soda, M., Mano, H., Togayachi, S., Miyoshi, H. et al. (2015). Epigenetic regulation of the nuclear-coded GCAT and SHMT2 genes confers human age-associated mitochondrial respiration defects. *Sci. Rep.* **5**, 10434.
- Hasty, P., Campisi, J., Hoeijmakers, J., van Steeg, H. and Vijg, J. (2003). Aging and genome maintenance: lessons from the mouse? *Science* **299**, 1355-1359.
- Helfand, S. L. and Rogina, B. (2003). Genetics of aging in the fruit fly, *Drosophila melanogaster*. *Annu. Rev. Genet.* **37**, 329-348.
- Henderson, K. A. and Gottschling, D. E. (2008). A mother's sacrifice: what is she keeping for herself? *Curr. Opin. Cell Biol.* **20**, 723-728.
- Henriques, C. M., Carneiro, M. C., Tenente, I. M., Jacinto, A. and Ferreira, M. G. (2013). Telomerase is required for zebrafish lifespan. *PLoS Genet.* **9**, e1003214.
- Herker, E., Jungwirth, H., Lehmann, K. A., Maldener, C., Frohlich, K.-U., Wissing, S., Buttner, S., Fehr, M., Sigrist, S. and Madeo, F. (2004). Chronological aging leads to apoptosis in yeast. *J. Cell Biol.* **164**, 501-507.
- Hertweck, M., Hoppe, T. and Baumeister, R. (2003). *C. elegans*, a model for aging with high-throughput capacity. *Exp. Gerontol.* **38**, 345-346.
- Hisaoka, K. K. and Firlit, C. F. (1962). Ovarian cycle and egg production in the zebrafish, *Brachydanio rerio*. *Copeia* **1962**, 788-792.
- Hwang, W. Y., Fu, Y., Reyon, D., Maeder, M. L., Tsai, S. Q., Sander, J. D., Peterson, R. T., Yeh, J.-R. and Jung, J. K. (2013). Efficient genome editing in zebrafish using a CRISPR-Cas system. *Nat. Biotechnol.* **31**, 227-229.
- Ingram, D. K. (1983). Toward the behavioral assessment of biological aging in the laboratory mouse: concepts, terminology, and objectives. *Exp. Aging Res.* **9**, 225-238.
- Jacobson, J., Lambert, A. J., Portero-Otín, M., Pamplona, R., Magwere, T., Miwa, S., Drieger, Y., Brand, M. D. and Partridge, L. (2010). Biomarkers of aging in *Drosophila*. *Aging Cell* **9**, 466-477.
- Jao, L.-E., Wente, S. R. and Chen, W. (2013). Efficient multiplex biallelic zebrafish genome editing using a CRISPR nuclease system. *Proc. Natl. Acad. Sci. USA* **110**, 13904-13909.
- Johnson, T. E. (2003). Advantages and disadvantages of *Caenorhabditis elegans* for aging research. *Exp. Gerontol.* **38**, 1329-1332.
- Johnson, T. E. and Hutchinson, E. W. (1993). Absence of strong heterosis for life span and other life history traits in *Caenorhabditis elegans*. *Genetics* **134**, 465-474.
- Johnson, D. K., Rinchik, E. M., Moustaid-Moussa, N., Miller, D. R., Williams, R. W., Michaud, E. J., Jablonski, M. M., Elberger, A., Hamre, K., Smeys, R. et al. (2005). Phenotype screening for genetically determined age-onset disorders and increased longevity in ENU-mutagenized mice. *Age* **27**, 75-90.
- Jorgensen, E. M. and Mango, S. E. (2002). The art and design of genetic screens: *Caenorhabditis elegans*. *Nat. Rev. Genet.* **3**, 356-369.
- Kaerberlein, M. (2010). Lessons on longevity from budding yeast. *Nature* **464**, 513-519.
- Kaerberlein, M., Powers, R. W., III, Steffen, K. K., Westman, E. A., Hu, D., Dang, N., Kerr, E. O., Kirkland, K. T., Fields, S. and Kennedy, B. K. (2005). Regulation of yeast replicative life span by TOR and Sch9 in response to nutrients. *Science* **310**, 1193-1196.
- Kaluff, A. V., Stewart, A. M. and Gerlai, R. (2014). Zebrafish as an emerging model for studying complex brain disorders. *Trends Pharmacol. Sci.* **35**, 63-75.
- Keane, M., Semeiks, J., Webb, A. E., Li, Y. I., Quesada, V., Craig, T., Madsen, L. B., van Dam, S., Brawand, D., Marques, P. I. et al. (2015). Insights into the evolution of longevity from the bowhead whale genome. *Cell Rep.* **10**, 112-122.
- Kenyon, C. J. (2010). The genetics of ageing. *Nature* **464**, 504-512.
- Kenyon, C., Chang, J., Gensch, E., Rudner, A. and Tabtiang, R. (1993). A *C. elegans* mutant that lives twice as long as wild type. *Nature* **366**, 461-464.
- Khurana, V. and Lindquist, S. (2010). Modelling neurodegeneration in *Saccharomyces cerevisiae*: why cook with baker's yeast? *Nat. Rev. Neurosci.* **11**, 436-449.
- Kikuchi, K. (2015). Dedifferentiation, transdifferentiation, and proliferation: mechanisms underlying cardiac muscle regeneration in zebrafish. *Curr. Pathobiol. Rep.* **3**, 81-88.
- Kim, S. K. (2007). Common aging pathways in worms, flies, mice and humans. *J. Exp. Biol.* **210**, 1607-1612.
- Kim, E. B., Fang, X., Fushan, A. A., Huang, Z., Lobanov, A. V., Han, L., Marino, S. M., Sun, X., Turanov, A. A., Yang, P. et al. (2011). Genome sequencing reveals insights into physiology and longevity of the naked mole rat. *Nature* **479**, 223-227.
- Kimmel, C. B., Ballard, W. W., Kimmel, S. R., Ullmann, B. and Schilling, T. F. (1995). Stages of embryonic development of the zebrafish. *Dev. Dyn.* **203**, 253-310.
- Kimura, K. D., Tissenbaum, H. A., Liu, Y. and Ruvkun, G. (1997). *daf-2*, an insulin receptor-like gene that regulates longevity and diapause in *Caenorhabditis elegans*. *Science* **277**, 942-946.
- Kirkwood, T. B. L. and Austad, S. N. (2000). Why do we age? *Nature* **408**, 233-238.
- Kirschner, J., Weber, D., Neuschl, C., Franke, A., Bottger, M., Zielke, L., Powalsky, E., Groth, M., Shagin, D., Petzold, A. et al. (2012). Mapping of quantitative trait loci controlling lifespan in the short-lived fish *Nothobranchius furzeri*—a new vertebrate model for age research. *Aging Cell* **11**, 252-261.
- Kishi, S. (2011). The search for evolutionary developmental origins of aging in zebrafish: a novel intersection of developmental and senescence biology in the zebrafish model system. *Birth Defects Res. C Embryo Today* **93**, 229-248.
- Kishi, S., Uchiyama, J., Baughman, A. M., Goto, T., Lin, M. C. and Tsai, S. B. (2003). The zebrafish as a vertebrate model of functional aging and very gradual senescence. *Exp. Gerontol.* **38**, 777-786.
- Kishi, S., Bayliss, P. E., Uchiyama, J., Koshimizu, E., Qi, J., Nanjappa, P., Imamura, S., Islam, A., Neuber, D., Amsterdam, A. et al. (2008). The identification of zebrafish mutants showing alterations in senescence-associated biomarkers. *PLoS Genet.* **4**, e1000152.
- Kitano, K. (2014). Structural mechanisms of human RecQ helicases WRN and BLM. *Front. Genet.* **5**, 366.
- Klass, M. R. (1977). Aging in the nematode *Caenorhabditis elegans*: major biological and environmental factors influencing life span. *Mech. Ageing Dev.* **6**, 413-429.
- Klass, M. R. (1983). A method for the isolation of longevity mutants in the nematode *Caenorhabditis elegans* and initial results. *Mech. Ageing Dev.* **22**, 279-286.
- Kurz, D. J., Decary, S., Hong, Y. and Erusalimsky, J. D. (2000). Senescence-associated (beta)-galactosidase reflects an increase in lysosomal mass during replicative ageing of human endothelial cells. *J. Cell Sci.* **113**, 3613-3622.
- Kutscher, L. M. and Shaham, S. (2014). Forward and reverse mutagenesis in *C. elegans*. *WormBook*, 1-26.
- Lamb, M. J. (1968). Temperature and lifespan in *Drosophila*. *Nature* **220**, 808-809.
- Lane, S. J., Frankino, W. A., Elekonich, M. M. and Roberts, S. P. (2014). The effects of age and lifetime flight behavior on flight capacity in *Drosophila melanogaster*. *J. Exp. Biol.* **217**, 1437-1443.
- Lawson, N. D. and Wolfe, S. A. (2011). Forward and reverse genetic approaches for the analysis of vertebrate development in the zebrafish. *Dev. Cell* **21**, 48-64.
- Lee, R. Y. N., Hench, J. and Ruvkun, G. (2001). Regulation of *C. elegans* DAF-16 and its human ortholog FKHRL1 by the *daf-2* insulin-like signaling pathway. *Curr. Biol.* **11**, 1950-1957.
- Lee, S. S., Lee, R. Y. N., Fraser, A. G., Kamath, R. S., Ahringer, J. and Ruvkun, G. (2003). A systematic RNAi screen identifies a critical role for mitochondria in *C. elegans* longevity. *Nat. Genet.* **33**, 40-48.
- Lewinska, A., Miedzak, B., Kulak, K., Molon, M. and Wnuk, M. (2014). Links between nucleolar activity, rDNA stability, aneuploidy and chronological aging in the yeast *Saccharomyces cerevisiae*. *Biogerontology* **15**, 289-316.
- Liao, C.-Y. and Kennedy, B. K. (2014). Mouse models and aging: longevity and progeria. *Curr. Top. Dev. Biol.* **109**, 249-285.
- Lieschke, G. J. and Currie, P. D. (2007). Animal models of human disease: zebrafish swim into view. *Nat. Rev. Genet.* **8**, 353-367.
- Lin, Y.-J., Seroude, L. and Benzer, S. (1998). Extended life-span and stress resistance in the *Drosophila* mutant *methuselah*. *Science* **282**, 943-946.
- Lithgow, G. J. (1996). Temperature, stress response and aging. *Rev. Clin. Gerontol.* **6**, 119-127.
- Lithgow, G. J., White, T. M., Melov, S. and Johnson, T. E. (1995). Thermotolerance and extended life-span conferred by single-gene mutations and induced by thermal stress. *Proc. Natl. Acad. Sci. USA* **92**, 7540-7544.
- Liu, R. K. and Walford, R. L. (1966). Increased growth and life-span with lowered ambient temperature in the annual fish, *Cynolebias adloffii*. *Nature* **212**, 1277-1278.
- Liu, T., Qi, H., Ma, L., Liu, Z., Fu, H., Zhu, W., Song, T., Yang, B. and Li, G. (2015). Resveratrol attenuates oxidative stress and extends lifespan in the annual fish *Nothobranchius Guentheri*. *Rejuvenat. Res.* **18**, 225-233.
- Longo, V. D., Gralla, E. B. and Valentine, J. S. (1996). Superoxide dismutase activity is essential for stationary phase survival in *Saccharomyces cerevisiae*: mitochondrial production of toxic oxygen species in vivo. *J. Biol. Chem.* **271**, 12275-12280.
- Longo, V. D., Ellerby, L. M., Bredesen, D. E., Valentine, J. S. and Gralla, E. B. (1997). Human Bcl-2 reverses survival defects in yeast lacking superoxide dismutase and delays death of wild-type yeast. *J. Cell Biol.* **137**, 1581-1588.
- Lopez-Otin, C., Blasco, M. A., Partridge, L., Serrano, M. and Kroemer, G. (2013). The hallmarks of aging. *Cell* **153**, 1194-1217.
- Mair, W., Goymer, P., Pletcher, S. D. and Partridge, L. (2003). Demography of dietary restriction and death in *Drosophila*. *Science* **301**, 1731-1733.
- Mair, W., Piper, M. D. W. and Partridge, L. (2005). Calories do not explain extension of life span by dietary restriction in *Drosophila*. *PLoS Biol.* **3**, e223.

- Marchal, J., Pifferi, F. and Aujard, F. (2013). Resveratrol in mammals: effects on aging biomarkers, age-related diseases, and life span. *Ann. N. Y. Acad. Sci.* **1290**, 67-73.
- Markofsky, J. and Matias, J. R. (1977). The effects of temperature and season of collection on the onset and duration of diapause in embryos of the annual fish *Nothobranchius guentheri*. *J. Exp. Zool.* **202**, 49-56.
- Markofsky, J. and Milstoc, M. (1979a). Aging changes in the liver of the male annual cyprinodont fish, *Nothobranchius guentheri*. *Exp. Gerontol.* **14**, 11-20.
- Markofsky, J. and Milstoc, M. (1979b). Histopathological observations of the kidney during aging of the male annual fish *Nothobranchius guentheri*. *Exp. Gerontol.* **14**, 149-155.
- Masoro, E. J. (1996). Handbook of the biology of aging, 4th edition - Schneider, E. L., Rowe, J. W. *Gerontologist* **36**, 828-830.
- Masoro, E. J. and Austad, S. N. (2006). *Handbook of the Biology of Aging*. Amsterdam; Boston: Elsevier Academic Press.
- Matthews, K. A., Kaufman, T. C. and Gelbart, W. M. (2005). Research resources for *Drosophila*: the expanding universe. *Nat. Rev. Genet.* **6**, 179-193.
- Micchelli, C. A. and Perrimon, N. (2006). Evidence that stem cells reside in the adult *Drosophila* midgut epithelium. *Nature* **439**, 475-479.
- Millburn, G. H., Crosby, M. A., Gramates, L. S., Tweedie, S. and the FlyBase Consortium (2016). FlyBase portals to human disease research using *Drosophila* models. *Dis. Model. Mech.* **9** (in press). doi:10.1242/dmm.023317.
- Miller, R. A., Buehner, G., Chang, Y., Harper, J. M., Sigler, R. and Smith-Wheelock, M. (2005). Methionine-deficient diet extends mouse lifespan, slows immune and lens aging, alters glucose, T4, IGF-I and insulin levels, and increases hepatocyte MIF levels and stress resistance. *Aging Cell* **4**, 119-125.
- Miller, C. T., Beleza, S., Pollen, A. A., Schluter, D., Kittles, R. A., Shriver, M. D. and Kingsley, D. M. (2007). cis-Regulatory changes in Kit ligand expression and parallel evolution of pigmentation in sticklebacks and humans. *Cell* **131**, 1179-1189.
- Moens, C. B., Donn, T. M., Wolf-Saxon, E. R. and Ma, T. P. (2008). Reverse genetics in zebrafish by TILLING. *Brief Funct. Genomic Proteomic* **7**, 454-459.
- Muller, H. J. (1927). Artificial transmutation of the gene. *Science* **66**, 84-87.
- Nordenstedt, H., White, D. L. and El-Serag, H. B. (2010). The changing pattern of epidemiology in hepatocellular carcinoma. *Dig. Liver Dis.* **42** Suppl. 3, S206-S214.
- Ocorr, K., Perrin, L., Lim, H.-Y., Qian, L., Wu, X. and Bodmer, R. (2007). Genetic control of heart function and aging in *Drosophila*. *Trends Cardiovasc. Med.* **17**, 177-182.
- Ogg, S., Paradis, S., Gottlieb, S., Patterson, G. I., Lee, L., Tissenbaum, H. A. and Ruvkun, G. (1997). The Fork head transcription factor DAF-16 transduces insulin-like metabolic and longevity signals in *C. elegans*. *Nature* **389**, 994-999.
- Ohlstein, B. and Spradling, A. (2006). The adult *Drosophila* posterior midgut is maintained by pluripotent stem cells. *Nature* **439**, 470-474.
- Patton, E. E. and Zon, L. I. (2001). The art and design of genetic screens: zebrafish. *Nat. Rev. Genet.* **2**, 956-966.
- Petzold, A., Reichwald, K., Groth, M., Taudien, S., Hartmann, N., Priebe, S., Shagin, D., Englert, C. and Platzer, M. (2013). The transcript catalogue of the short-lived fish *Nothobranchius furzeri* provides insights into age-dependent changes of mRNA levels. *BMC Genomics* **14**, 185.
- Piper, M. D. W., Partridge, L., Raubenheimer, D. and Simpson, S. J. (2011). Dietary restriction and aging: a unifying perspective. *Cell Metab.* **14**, 154-160.
- Podrabsky, J. E. (1999). Husbandry of the annual killifish *Austrofundulus limnaeus* with special emphasis on the collection and rearing of embryos. *Environ. Biol. Fishes* **54**, 421-431.
- Podrabsky, J. E., Garrett, I. D. F. and Kohl, Z. F. (2010). Alternative developmental pathways associated with diapause regulated by temperature and maternal influences in embryos of the annual killifish *Austrofundulus limnaeus*. *J. Exp. Biol.* **213**, 3280-3288.
- Polacik, M. and Reichard, M. (2011). Asymmetric reproductive isolation between two sympatric annual killifish with extremely short lifespans. *PLoS ONE* **6**, e22684.
- Pompei, F., Polkanov, M. and Wilson, R. (2001). Age distribution of cancer in mice: the incidence turnover at old age. *Toxicol. Ind. Health* **17**, 7-16.
- Poss, K. D., Wilson, L. G. and Keating, M. T. (2002). Heart regeneration in zebrafish. *Science* **298**, 2188-2190.
- Poss, K. D., Keating, M. T. and Nechiporuk, A. (2003). Tales of regeneration in zebrafish. *Dev. Dyn.* **226**, 202-210.
- Powers, R. W., III, Kaerberlein, M., Caldwell, S. D., Kennedy, B. K. and Fields, S. (2006). Extension of chronological life span in yeast by decreased TOR pathway signaling. *Genes Dev.* **20**, 174-184.
- Promislow, D. E., Tatar, M., Khazaeli, A. A. and Curtsinger, J. W. (1996). Age-specific patterns of genetic variance in *Drosophila melanogaster*. I. Mortality. *Genetics* **143**, 839-848.
- Reichard, M., Polacik, M. and Sedlacek, O. (2009). Distribution, colour polymorphism and habitat use of the African killifish *Nothobranchius furzeri*, the vertebrate with the shortest life span. *J. Fish Biol.* **74**, 198-212.
- Reichwald, K., Lauber, C., Nanda, I., Kirschner, J., Hartmann, N., Schories, S., Gausmann, U., Taudien, S., Schilhabel, M. B., Szafranski, K. et al. (2009). High tandem repeat content in the genome of the short-lived annual fish *Nothobranchius furzeri*: a new vertebrate model for aging research. *Genome Biol.* **10**, R16.
- Reichwald, K., Petzold, A., Koch, P., Downie, B. R., Hartmann, N., Pietsch, S., Baumgart, M., Chalopin, D., Felder, M., Bens, M. et al. (2015). Insights into sex chromosome evolution and aging from the genome of a short-lived fish. *Cell* **163**, 1527-1538.
- Reznick, D., Bryant, M. and Holmes, D. (2006). The evolution of senescence and post-reproductive lifespan in guppies (*Poecilia reticulata*). *PLoS Biol.* **4**, e7.
- Riddle, D. L., Swanson, M. M. and Albert, P. S. (1981). Interacting genes in nematode dauer larva formation. *Nature* **290**, 668-671.
- Ruzicka, L., Bradford, Y. M., Frazer, K., Howe, D. G., Paddock, H., Ramachandran, S., Singer, A., Toro, S., Van Slyke, C. E., Eagle, A. E. et al. (2015). ZFIN, The zebrafish model organism database: updates and new directions. *Genesis* **53**, 498-509.
- Sakaue-Sawano, A., Kurokawa, H., Morimura, T., Hanyu, A., Hama, H., Osawa, H., Kashiwagi, S., Fukami, K., Miyata, T., Miyoshi, H. et al. (2008). Visualizing spatiotemporal dynamics of multicellular cell-cycle progression. *Cell* **132**, 487-498.
- Schmued, L. C. and Hopkins, K. J. (2000). Fluoro-Jade B: a high affinity fluorescent marker for the localization of neuronal degeneration. *Brain Res.* **874**, 123-130.
- Sebastiani, P., Riva, A., Montano, M., Pham, P., Torkamani, A., Scherba, E., Benson, G., Milton, J. N., Baldwin, C. T., Andersen, S. et al. (2011). Whole genome sequences of a male and female supercentenarian, ages greater than 114 years. *Front. Genet.* **2**, 90.
- Sinclair, D. A. and Guarente, L. (1997). Extrachromosomal rDNA circles—a cause of aging in yeast. *Cell* **91**, 1033-1042.
- Skorupa, D. A., Dervisevendic, A., Zwiener, J. and Pletcher, S. D. (2008). Dietary composition specifies consumption, obesity, and lifespan in *Drosophila melanogaster*. *Aging Cell* **7**, 478-490.
- Solon-Biet, S. M., Mitchell, S. J., Coogan, S. C. P., Cogger, V. C., Gokarn, R., McMahon, A. C., Raubenheimer, D., de Cabo, R., Simpson, S. J. and Le Couteur, D. G. (2015). Dietary protein to carbohydrate ratio and caloric restriction: comparing metabolic outcomes in mice. *Cell Rep.* **11**, 1529-1534.
- Stanfel, M. N., Shamieh, L. S., Kaerberlein, M. and Kennedy, B. K. (2009). The TOR pathway comes of age. *Biochim. Biophys. Acta* **1790**, 1067-1074.
- Sung, Y. H., Baek, I.-J., Kim, D. H., Jeon, J., Lee, J., Lee, K., Jeong, D., Kim, J.-S. and Lee, H.-W. (2013). Knockout mice created by TALEN-mediated gene targeting. *Nat. Biotechnol.* **31**, 23-24.
- Tacutu, R., Craig, T., Budovsky, A., Wuttke, D., Lehmann, G., Taranukha, D., Costa, J., Fraifeld, V. E. and de Magalhães, J. P. (2013). Human Ageing Genomic Resources: integrated databases and tools for the biology and genetics of ageing. *Nucleic Acids Res.* **41**, D1027-D1033.
- Taormina, G. and Mirisola, M. G. (2014). Calorie restriction in mammals and simple model organisms. *Biomed. Res. Int.* **2014**, 308690.
- Tatar, M., Promislow, D. E., Khazaeli, A. A. and Curtsinger, J. W. (1996). Age-specific patterns of genetic variance in *Drosophila melanogaster*. II. Fecundity and its genetic covariance with age-specific mortality. *Genetics* **143**, 849-858.
- Tatarenkov, A. and Avise, J. C. (2007). Rapid concerted evolution in animal mitochondrial DNA. *Proc. Biol. Sci.* **274**, 1795-1798.
- Tauchi, H. and Sato, T. (1968). Age changes in size and number of mitochondria of human hepatic cells. *J. Gerontol.* **23**, 454-461.
- Terzibasi, E., Valenzano, D. R. and Cellerino, A. (2007). The short-lived fish *Nothobranchius furzeri* as a new model system for aging studies. *Exp. Gerontol.* **42**, 81-89.
- Terzibasi, E., Valenzano, D. R., Benedetti, M., Roncaglia, P., Cattaneo, A., Domenici, L. and Cellerino, A. (2008). Large differences in aging phenotype between strains of the short-lived annual fish *Nothobranchius furzeri*. *PLoS ONE* **3**, e3866.
- Terzibasi, E., Lefrançois, C., Domenici, P., Hartmann, N., Graf, M. and Cellerino, A. (2009). Effects of dietary restriction on mortality and age-related phenotypes in the short-lived fish *Nothobranchius furzeri*. *Aging Cell* **8**, 88-99.
- Tissenbaum, H. A. (2015). Using *C. elegans* for aging research. *Invertebr. Reprod. Dev.* **59**, 59-63.
- Tissenbaum, H. A. and Ruvkun, G. (1998). An insulin-like signaling pathway affects both longevity and reproduction in *Caenorhabditis elegans*. *Genetics* **148**, 703-717.
- Tozzini, E. T., Baumgart, M., Battistoni, G. and Cellerino, A. (2012). Adult neurogenesis in the short-lived teleost *Nothobranchius furzeri*: localization of neurogenic niches, molecular characterization and effects of aging. *Aging Cell* **11**, 241-251.
- Untergasser, G., Gander, R., Rumpold, H., Heinrich, E., Plas, E. and Berger, P. (2003). TGF-beta cytokines increase senescence-associated beta-galactosidase activity in human prostate basal cells by supporting differentiation processes, but not cellular senescence. *Exp. Gerontol.* **38**, 1179-1188.
- Valdesalici, S. and Cellerino, A. (2003). Extremely short lifespan in the annual fish *Nothobranchius furzeri*. *Proc. Biol. Sci.* **270** Suppl. 2, S189-S191.
- Valenzano, D. R. and Cellerino, A. (2006). Resveratrol and the pharmacology of aging: a new vertebrate model to validate an old molecule. *Cell Cycle* **5**, 1027-1032.

- Valenzano, D. R., Terzibasi, E., Cattaneo, A., Domenici, L. and Cellerino, A.** (2006a). Temperature affects longevity and age-related locomotor and cognitive decay in the short-lived fish *Nothobranchius furzeri*. *Aging Cell* **5**, 275-278.
- Valenzano, D. R., Terzibasi, E., Genade, T., Cattaneo, A., Domenici, L. and Cellerino, A.** (2006b). Resveratrol prolongs lifespan and retards the onset of age-related markers in a short-lived vertebrate. *Curr. Biol.* **16**, 296-300.
- Valenzano, D. R., Kirschner, J., Kamber, R. A., Zhang, E., Weber, D., Cellerino, A., Englert, C., Platzer, M., Reichwald, K. and Brunet, A.** (2009). Mapping loci associated with tail color and sex determination in the short-lived fish *Nothobranchius furzeri*. *Genetics* **183**, 1385-1395.
- Valenzano, D. R., Sharp, S. and Brunet, A.** (2011). Transposon-mediated transgenesis in the short-lived African killifish *Nothobranchius furzeri*, a vertebrate model for aging. *G3* **1**, 531-538.
- Valenzano, D. R., Benayoun, B. A., Singh, P. P., Zhang, E., Etter, P. D., Hu, C.-K., Clément-Ziza, M., Willemsen, D., Cui, R., Harel, I. et al.** (2015). The African turquoise killifish genome provides insights into evolution and genetic architecture of lifespan. *Cell* **163**, 1539-1554.
- Van Houcke, J., De Groef, L., Dekeyster, E. and Moons, L.** (2015). The zebrafish as a gerontology model in nervous system aging, disease, and repair. *Ageing Res. Rev.* **24**, 358-368.
- Vanhooren, V. and Libert, C.** (2013). The mouse as a model organism in aging research: usefulness, pitfalls and possibilities. *Ageing Res. Rev.* **12**, 8-21.
- Veith, S. and Mangerich, A.** (2015). RecQ helicases and PARP1 team up in maintaining genome integrity. *Ageing Res. Rev.* **23**, 12-28.
- Veldman, M. B. and Lin, S.** (2008). Zebrafish as a developmental model organism for pediatric research. *Pediatr. Res.* **64**, 470-476.
- Verduyck, M., Vignaud, H., Bynens, T., Van den Brande, J., Franssens, V., Cullin, C. and Winderickx, J.** (2016). Yeast as a model for Alzheimer's disease: latest studies and advanced strategies. *Methods Mol. Biol.* **1303**, 197-215.
- Vermulst, M., Wanagat, J., Kujoth, G. C., Bielas, J. H., Rabinovitch, P. S., Prolla, T. A. and Loeb, L. A.** (2008). DNA deletions and clonal mutations drive premature aging in mitochondrial mutator mice. *Nat. Genet.* **40**, 392-394.
- Walker, G. A., Walker, D. W. and Lithgow, G. J.** (1998). Genes that determine both thermotolerance and rate of aging in *Caenorhabditis elegans*. *Stress Life* **851**, 444-449.
- Wallenfang, M. R., Nayak, R. and DiNardo, S.** (2006). Dynamics of the male germline stem cell population during aging of *Drosophila melanogaster*. *Aging Cell* **5**, 297-304.
- Walter, M. F., Biessmann, M. R., Benitez, C., Torok, T., Mason, J. M. and Biessmann, H.** (2007). Effects of telomere length in *Drosophila melanogaster* on life span, fecundity, and fertility. *Chromosoma* **116**, 41-51.
- Wang, K., Huang, Z., Zhao, L., Liu, W., Chen, X., Meng, P., Lin, Q., Chi, Y., Xu, M., Ma, N. et al.** (2012). Large-scale forward genetic screening analysis of development of hematopoiesis in zebrafish. *J. Genet. Genomics* **39**, 473-480.
- Wang, H., Yang, H., Shivalila, C. S., Dawlaty, M. M., Cheng, A. W., Zhang, F. and Jaenisch, R.** (2013). One-step generation of mice carrying mutations in multiple genes by CRISPR/Cas-mediated genome engineering. *Cell* **153**, 910-918.
- Wei, M., Fabrizio, P., Madia, F., Hu, J., Ge, H., Li, L. M. and Longo, V. D.** (2009). Tor1/Sch9-regulated carbon source substitution is as effective as calorie restriction in life span extension. *PLoS Genet.* **5**, e1000467.
- Wendler, S., Hartmann, N., Hoppe, B. and Englert, C.** (2015). Age-dependent decline in fin regenerative capacity in the short-lived fish *Nothobranchius furzeri*. *Aging Cell* **14**, 857-866.
- Westerfield, M.** (2000). *The Zebrafish Book. A Guide for the Laboratory Use of Zebrafish (Danio rerio)*. Eugene, Oregon: University of Oregon Press.
- WHO** (2011). Global health and ageing. National Institute on Aging.
- Willcox, D. C., Willcox, B. J., Hsueh, W.-C. and Suzuki, M.** (2006). Genetic determinants of exceptional human longevity: insights from the Okinawa Centenarian Study. *Age* **28**, 313-332.
- Williams, G. C.** (1957). Pleiotropy, natural selection, and the evolution of senescence. *Evolution* **11**, 398-411.
- Wong, A., Boutis, P. and Hekimi, S.** (1995). Mutations in the *clk-1* gene of *Caenorhabditis elegans* affect developmental and behavioral timing. *Genetics* **139**, 1247-1259.
- Woodhead, A. D. and Pond, V.** (1984). Aging changes in the optic tectum of the guppy *Poecilia (Lebistes) reticulatus*. *Exp. Gerontol.* **19**, 305-311.
- Woodhead, A. D., Pond, V. and Dailey, K.** (1983). Aging changes in the kidneys of two poeciliid fishes, the guppy *Poecilia reticulatus* and the Amazon molly *P. formosa*. *Exp. Gerontol.* **18**, 211-221.
- Wourms, J. P.** (1972). The developmental biology of annual fishes. III. Pre-embryonic and embryonic diapause of variable duration in the eggs of annual fishes. *J. Exp. Zool.* **182**, 389-414.
- Yang, Y. and Wilson, D. L.** (2000). Isolating aging mutants: a novel method yields three strains of the nematode *Caenorhabditis elegans* with extended life spans. *Mech. Ageing Dev.* **113**, 101-116.
- Yang, H., Wang, H. and Jaenisch, R.** (2014). Generating genetically modified mice using CRISPR/Cas-mediated genome engineering. *Nat. Protoc.* **9**, 1956-1968.
- Yanos, M. E., Bennett, C. F. and Kaerberlein, M.** (2012). Genome-wide RNAi longevity screens in *Caenorhabditis elegans*. *Curr. Genomics* **13**, 508-518.
- Yegorov, Y. E., Akimov, S. S., Hass, R., Zelenin, A. V. and Prudovsky, I. A.** (1998). Endogenous beta-galactosidase activity in continuously nonproliferating cells. *Exp. Cell Res.* **243**, 207-211.
- Yen, T.-C., Chen, Y.-S., King, K.-L., Yeh, S.-H. and Wei, Y.-H.** (1989). Liver mitochondrial respiratory functions decline with age. *Biochem. Biophys. Res. Commun.* **165**, 994-1003.
- Yu, X. and Li, G.** (2012). Effects of resveratrol on longevity, cognitive ability and aging-related histological markers in the annual fish *Nothobranchius guentheri*. *Exp. Gerontol.* **47**, 940-949.
- Yuan, R., Tsaih, S.-W., Petkova, S. B., de Evsikova, C. M., Xing, S., Marion, M. A., Bogue, M. A., Mills, K. D., Peters, L. L., Bult, C. J. et al.** (2009). Aging in inbred strains of mice: study design and interim report on median lifespans and circulating IGF1 levels. *Aging Cell* **8**, 277-287.
- Yuan, R., Peters, L. L. and Paigen, B.** (2011). Mice as a mammalian model for research on the genetics of aging. *ILAR J.* **52**, 4-15.
- Yui, R., Ohno, Y. and Matsuura, E. T.** (2003). Accumulation of deleted mitochondrial DNA in aging *Drosophila melanogaster*. *Genes Genet. Syst.* **78**, 245-251.
- Zahn, J. M., Poosala, S., Owen, A. B., Ingram, D. K., Lustig, A., Carter, A., Weeraratna, A. T., Taub, D. D., Gorospe, M., Mazan-Mamczarz, K. et al.** (2007). AGEMAP: a gene expression database for aging in mice. *PLoS Genet.* **3**, e201.
- Zhu, L., Hathcock, K. S., Hande, P., Lansdorp, P. M., Seldin, M. F. and Hodes, R. J.** (1998). Telomere length regulation in mice is linked to a novel chromosome locus. *Proc. Natl. Acad. Sci. USA* **95**, 8648-8653.
- Zimmerman, J. A., Malloy, V., Krajcik, R. and Orentreich, N.** (2003). Nutritional control of aging. *Exp. Gerontol.* **38**, 47-52.